

TOPICAL REVIEW • OPEN ACCESS

From legacy contamination to watershed systems science: a review of scientific insights and technologies developed through DOE-supported research in water and energy security

To cite this article: Dipankar Dwivedi *et al* 2022 *Environ. Res. Lett.* **17** 043004

View the [article online](#) for updates and enhancements.

You may also like

- [Kupka–Smale diffeomorphisms at the boundary of uniform hyperbolicity: a model](#)
Renaud Leplaideur and Isabel Lugão Rios
- [Mountaintop mining legacies constrain ecological, hydrological and biogeochemical recovery trajectories](#)
Matthew R V Ross, Fabian Nippgen, Brian L McGlynn *et al.*
- [QMCPACK: an open source *ab initio* quantum Monte Carlo package for the electronic structure of atoms, molecules and solids](#)
Jeongnim Kim, Andrew D Baczewski, Todd D Beaudet *et al.*

ENVIRONMENTAL RESEARCH
LETTERS

TOPICAL REVIEW

OPEN ACCESS

RECEIVED
30 March 2021REVISED
6 February 2022ACCEPTED FOR PUBLICATION
1 March 2022PUBLISHED
18 March 2022

Original Content from
this work may be used
under the terms of the
[Creative Commons
Attribution 4.0 licence](#).

Any further distribution
of this work must
maintain attribution to
the author(s) and the title
of the work, journal
citation and DOI.



From legacy contamination to watershed systems science: a review of scientific insights and technologies developed through DOE-supported research in water and energy security

Dipankar Dwivedi^{1,*}, Carl I Steefel¹, Bhavna Arora¹, Jill Banfield², John Bargar³, Maxim I Boyanov⁴, Scott C Brooks⁵, Xingyuan Chen⁶, Susan S Hubbard¹, Dan Kaplan⁷, Kenneth M Kemner⁴, Peter S Nico¹, Edward J O'Loughlin⁴, Eric M Pierce⁵, Scott L Painter⁵, Timothy D Scheibe⁶, Haruko M Wainwright^{1,8}, Kenneth H Williams¹ and Mavrik Zavarin⁹

¹ Lawrence Berkeley National Laboratory, Berkeley, CA, United States of America

² University of California, Berkeley, CA, United States of America

³ SLAC National Accelerator Laboratory, Menlo Park, CA, United States of America

⁴ Argonne National Laboratory, Lemont, IL, United States of America

⁵ Oak Ridge National Laboratory, Oak Ridge, TN, United States of America

⁶ Pacific Northwest National Laboratory, Richland, WA, United States of America

⁷ Savannah River Ecology Laboratory, Jackson, SC, United States of America

⁸ Massachusetts Institute of Technology, Cambridge, MA, United States of America

⁹ Lawrence Livermore National Laboratory, Livermore, CA, United States of America

* Author to whom any correspondence should be addressed.

E-mail: DDwivedi@lbl.gov

Keywords: contaminant, groundwater, critical zone, reactive transport models, redox, hot spots and hot moments

Abstract

Water resources, including groundwater and prominent rivers worldwide, are under duress because of excessive contaminant and nutrient loads. To help mitigate this problem, the United States Department of Energy (DOE) has supported research since the late 1980s to improve our fundamental knowledge of processes that could be used to help clean up challenging subsurface problems. Problems of interest have included subsurface radioactive waste, heavy metals, and metalloids (e.g. uranium, mercury, arsenic). Research efforts have provided insights into detailed groundwater biogeochemical process coupling and the resulting geochemical exports of metals and nutrients to surrounding environments. Recently, an increased focus has been placed on constraining the exchanges and fates of carbon and nitrogen within and across bedrock to canopy compartments of a watershed and in river–floodplain settings, because of their important role in driving biogeochemical interactions with contaminants and the potential of increased fluxes under changing precipitation regimes, including extreme events. While reviewing the extensive research that has been conducted at DOE's representative sites and testbeds (such as the Oyster Site in Virginia, Savannah River Site in South Carolina, Oak Ridge Reservation in Tennessee, Hanford in Washington, Nevada National Security Site in Nevada, Riverton in Wyoming, and Rifle and East River in Colorado), this review paper explores the nature and distribution of contaminants in the surface and shallow subsurface (i.e. the critical zone) and their interactions with carbon and nitrogen dynamics. We also describe state-of-the-art, scale-aware characterization approaches and models developed to predict contaminant fate and transport. The models take advantage of DOE leadership-class high-performance computers and are beginning to incorporate artificial intelligence approaches to tackle the extreme diversity of hydro-biogeochemical processes and measurements. Recognizing that the insights and capability developments are potentially transferable to many other sites, we also explore the scientific implications of these advances and recommend future research directions.

1. Introduction

Water security is critical for food and energy production, economic development, and national security. Yet water security is under severe duress globally because of climate change, growing population, and human activities (Alley *et al* 2002, Rodell *et al* 2009, Heathwaite 2010, Famiglietti 2014). A recent study projected that more than 65% of the human population (~5 billion people) live in water-insecure regions (Vörösmarty *et al* 2010). Exacerbating the problem, the demand for freshwater (<3% of all water on Earth) is increasing, challenging our ability to meet food and energy needs globally. Although future technologies may allow us to increase clean water supplies, it is imperative to protect the available freshwater resources from numerous threats, such as a range of chemicals, including metals, metalloids, radionuclides, and nutrients that reduce usable supply.

Increasing contamination of freshwater poses a serious problem to both surface water and groundwater (e.g. Varol and Şen 2012, Brender *et al* 2013, Dwivedi *et al* 2013, 2016b), particularly because most of the world's prominent rivers, supporting several million people, are under threat and experiencing contamination from a variety of chemicals, including uranium (U), chromium (Cr), arsenic (As) and excessive amounts of nutrients (e.g. Brown and Halweil 1998, Dodds 2006, Paddison 2016, Tripathi *et al* 2016, Gross 2017, Lu *et al* 2018). To understand the behavior and reactivity of a plethora of contaminants and nutrients, the United State Department of Energy (DOE) has funded extensive research across several sites in the United States of America (USA) for several decades now (www.energy.gov/). Below, we provide how scientific progress has evolved through these efforts (figure 1).

The DOE's mission is 'to ensure America's security and prosperity by addressing its energy, environmental and nuclear challenges through transformative science and technology solutions.' The DOE was created in 1977, succeeding various energy-related programs previously dispersed throughout various federal agencies, including the Manhattan Project effort to develop the atomic bomb during World War II. Over fifty years of nuclear weapons production, testing, and energy research generated a vast volume of legacy contamination and created contaminated soil and water across the USA. The DOE has supported research and cleanup associated with several challenging subsurface problems since the late 1980s, such as treating radioactive waste (e.g. U, plutonium (Pu)) and heavy metals and metalloids (e.g. Cr, mercury (Hg), As) in the subsurface. Notably, since its inception in 1989, the DOE's Environmental Management (EM) and Environmental Remediation (ER) programs have been responsible for the restoration of

as many as 107 sites across the country—an area equal to Rhode Island and Delaware combined (figure 2). Within the DOE's Office of Science, the Biological and Environmental Research (BER) program has led the fundamental science research associated with water and energy security at the watershed, continental, and global scales through various projects and programs. The DOE-BER, its predecessors, and other related programs have significantly contributed to the progress of environmental sciences, setting the stage for this paper's thematic organization (figure 1).

Next, we briefly describe the history of various DOE programs that have supported water-security-research. The DOE-BER pioneered genomics, system biology, and biotechnology research through the Human Genome Project in 1990. In the late eighties, the Subsurface Science program initiated biological and co-contaminant geochemistry research through its deep microbiology and later bacterial transport projects. The Natural and Accelerated BIoRemediation (NABIR) program subsequently pioneered the 'bio' part of biogeochemistry (e.g. metal-reducing bacteria, in-silico modeling). The Subsurface Biogeochemical Research (SBR) program then continued that linkage, emphasizing the interplay between transport and biogeochemistry. Within the DOE's Office of Science, the Advanced Scientific Computing Research (ASCR) brings together researchers from various fields to tackle some of the most challenging scientific problems, leveraging DOE's leadership-class supercomputers and developing high-end computational science. In the environmental sciences, the ASCR's Scientific Discovery Through Advanced Computing (SciDAC) program has boosted subsurface research like multiscale science and reactive transport modeling. We use 'DOE research' and 'DOE-supported science' interchangeably to broadly reflect scientific advances pioneered through BER-SBR, their predecessors, and other related programs (e.g. EM, ASCR). This review paper follows the evolution of DOE research in this context, focusing primarily on the subsurface environmental sciences from the nineties to recent times.

Research efforts during the 1990s and early 2000s led to significant developments in our understanding of microbial communities, metals geochemistry, and co-contaminant interactions. These included the development of novel techniques for sampling and cultivating subsurface microbial communities (Ghiorse and Wobber 1989, Phelps *et al* 1989), identification of colloids as an important mechanism for mobilizing low solubility metals and radionuclides (e.g. Kersting *et al* 1999, Santschi *et al* 2002, Suzuki *et al* 2002), understanding of the influence of natural organic matter (NOM) on metal speciation and sorption on mineral surfaces (Gu *et al* 1994, 1995), application of synchrotron-based x-ray absorption spectroscopies to elucidate the nature of metal and





Figure 1. The spiral shows the progression of DOE-supported science that evolved in both scale and complexity. Significant scientific progress has been achieved from fundamental subsurface microbiology to geochemistry and biogeochemistry to hydro-biogeochemistry and now to eco-hydro-biogeochemistry over the past two decades.

organic matter complexes on mineral surfaces (e.g. Bargar *et al* 2000) and bacterial cells (Kelly *et al* 2002, Boyanov *et al* 2003), computational approaches to understanding the fundamental properties of natural minerals as metal sorbents (e.g. Zachara *et al* (1995), Felmy and Rustad (1998), Steefel *et al* (2003)), examination of microbial impacts on metal immobilization (e.g. Fredrickson *et al* 2000, Labrenz *et al* 2000, Kemner *et al* 2004), and implementation of mechanistic processes in reactive transport codes to simulate transport behavior (e.g. Steefel *et al* 2005).

At the same time, it was realized that natural and accelerated bioremediation held great potential for solving a wide spectrum of contamination problems (Lovley and Phillips 1992). These research efforts led to significant advances in natural and active stabilization of metals in the subsurface (Bender *et al* 2000, Lovley 2003). Coupled hydrological and biogeochemical processes such as reactive, colloidal,

and advective–dispersive transport were investigated within the view of biological availability, transformation, and movement of radionuclides and metals (Wang *et al* 2003). Some notable examples include the abiotic (O’Loughlin *et al* 2003a) and microbially mediated reductive precipitation and stabilization of U, iron (Fe) reduction, Pu surface-mediated reduction (Powell *et al* 2005), and the influence of NOM (McCarthy *et al* 1998), and other competing processes that can limit metal immobilization in the subsurface (Brooks *et al* 2003). Subsequent studies (e.g. Zheng *et al* (2003), Wan *et al* (2005), Tokunaga *et al* (2008)) questioned whether the microbially mediated redox changes resulting in radionuclide immobilization were sustainable.

Motivated by the potential of natural and accelerated bioremediation as an effective cleanup strategy for DOE, the Oyster Virginia Subsurface Bacterial Transport Project was initiated in 1999



as the first large, team-based, multidisciplinary, multi-institutional, DOE-supported scientific project that was specifically tied to a field research testbed. Through iterative integration of information gained through lab experiments, field characterization, field experiments, and numerical modeling associated with the seminal South Oyster Site, the project greatly advanced a predictive understanding of bacterial transport in physically, chemically, and biologically heterogeneous aquifers. The project led to several scientific firsts, including the first:

- (a) Estimation of shallow subsurface moisture content (Hubbard *et al* 2001), hydraulic conductivity (Hubbard *et al* 2001), and Fe geochemistry (Chen *et al* 2004) using high-resolution geophysical data,
- (b) Use of geophysically-obtained estimates, field experimental datasets, and numerical models to document how even mild heterogeneity influenced transport (Hubbard *et al* 2001, Scheibe *et al* 2006),
- (c) Documentation of scale effects and relative controls of physical and geochemical heterogeneity on bacterial transport using numerical models (Scheibe and Chien 2003, Scheibe *et al* 2006), and
- (d) Development of new methods for microbial enumeration and strain selection (Fuller *et al* 2000, DeFlaun *et al* 2001, Johnson *et al* 2001) and new insights regarding microbial attachment and transport (Dong *et al* 2002a, 2002b, Mailloux *et al* 2003).

As summarized by Scheibe *et al* (2011), not only has the body of literature resulting from the Oyster Site research been widely cited, but the project has served as a model for subsequent DOE-supported team-based research using data from highly-instrumented field testbeds with model-guided experimental design (Scheibe *et al* 2001). Many such representative DOE sites and testbeds are described below (section 3), together with advanced characterization and simulation modeling approaches to advance predictive understanding of complex

hydro-biogeochemical phenomena and water security research.

In the late 2000s, much emphasis was placed on linking the interplay between transport and biogeochemistry (e.g. diffusion-limited mass transfer, kinetic sorption-desorption). Indeed, this was the time when a paradigm shift ensued in multiscale, multiphysics science, and reactive transport modeling (e.g. White and Oostrom 2000, 2003, Pruess 2004, Steefel *et al* 2005, Hammond *et al* 2007). These advances subsequently paved the way for fully exploiting the DOE's unique computational capabilities in high-performance computing. In unison, subsurface environmental simulation capabilities were enhanced by explicitly representing multiphase, multicomponent process dynamics, and by code parallelization.

Up until the past decade, the investigations of small-scale (molecular to millimeter) processes had been a primary focus. This small-scale process understanding was used to upscale and predict behavior at the field scale by deriving scaling laws and mathematical models. Although this reductionist (bottom-up) approach was remarkable in advancing basic research, it was inadequate for addressing scaling behavior in the presence of a range of complex as well as coupled processes, nonlinearity, and a wide range of landscape heterogeneity. Consequently, large uncertainties existed in predictions that lacked a holistic, unifying approach to capture large-scale responses. There was a need to link small-scale process understanding with the larger-scale responses systematically (i.e. treating several ecosystem subsystems, components, and compartments, from the bedrock to the canopy, as a complex system). This perspective aims to comprehensively couple and model key processes across the critical zone (National Research Council 2001), and to link these processes to deep hydrological flow paths and atmospheric circulation systems.

To address this need, at the beginning of the last decade, researchers associated with DOE fate and transport challenges recommended new approaches that melded select strengths of mechanistic, bottom-up approaches that were prevalent at the time with top-down approaches, with an aim to improve characterization and prediction of complex hydro-biogeochemical behavior across scales (DOE-Complexity 2009). New emphasis was placed on actively linking sub-compartments of the Earth system, including atmospheric and biospheric processes, from the bedrock to the canopy. Multiple aspects of the system, such as hydrologic and carbon (C) cycles as well as microbiological and geochemical processes, were explicitly represented, linked, and allowed to interact with each other. This approach encouraged researchers to recognize scale transitions in a hierarchical subsurface system, link local and regional chemical fluxes, and investigate collective system behavior.

Subsequently, a more holistic system science approach evolved through active contaminant remediation-type studies. The system science approach tackles questions such as how watersheds respond to changes over the long term (e.g. Hubbard *et al* 2018). These include developing insights into the surface–subsurface hydrology; groundwater–surface water interactions; C, nutrient, and trace element transformations; ecosystem disturbances and resilience; impacts on earth systems, development of Earth System Models, and pioneering interdisciplinary community science (e.g. Stegen and Goldman 2018, Arora *et al* 2019b, Hubbard *et al* 2020).

Here we summarize the important findings developed from several decades of research work at DOE sites that have contributed to our understanding of contaminants and nutrient cycling and their transport in the environment. These DOE sites have been heavily tracked, monitored, and investigated for legacy contamination and have subsequently served as testbeds for exploring the exchange of materials between terrestrial and aquatic ecosystems—and how this exchange influences element export at river-basin scales. Therefore, these sites and their associated scientific discoveries provide an opportunity for us to review the nature and distribution of legacy contamination and their coupled interactions with C, nitrogen (N), and sulfur (S) dynamics, as well as the timescales of biogeochemical exports. Finally, with this article, we expect to raise awareness of the water-security problems arising from the dispersal of a range of chemicals in the environment and to provide a rich body of literature to address these problems.

2. An outline of DOE-supported science

A central focus of the DOE-supported science has been identifying the dominant biogeochemical processes controlling the migration of metals and radionuclides in the environment. Given the interrelationships between metals' behaviors and the presence of other elements (e.g. C, N, Fe, S), investigations into the geochemistry of metals and radionuclides have necessarily evolved to include detailed studies into the behavior of a broader range of elements that impact water quality (e.g. C, N). In the past decade, detailed mechanistic studies of metals geochemistry have included both laboratory and field investigations. Particular focus was initially placed on elements such as U (e.g. Rifle, Hanford, Oak Ridge), Hg (e.g. Oak Ridge), Pu (e.g. Savannah River Site, Nevada National Security Site), and metal co-contaminants (e.g. Rifle, Hanford), which later expanded to a broader set of biogeochemically complex elements, such as C and N, that are relevant to the ecosystem's health and water security at a national scale. Although it is not possible to capture all the scientific progress made at the DOE offices

and associated sites in the past several decades, our goal is to provide a broad understanding from all the contaminated sites in the past that lead to current ecosystem science.

We first describe the salient features of the DOE's representative sites used to achieve important successes in subsurface environmental sciences over the past two decades in section 3. Section 4 chronicles scientific progress in subsurface microbiology, geochemistry and biogeochemistry, and hydro-biogeochemistry over the past two decades. We primarily discuss the water security issues concerning the fate and transport of legacy contaminants, critical elements, and nutrients at representative DOE sites.

We then describe the capability development achieved while making many scientific advancements in section 5. We choose Hg as a use case to describe watershed biogeochemical processes holistically in section 6. Hg is emblematic in its complexity, unique to DOE challenges, and is particularly sensitive to extreme events that are increasingly expected in the future.

Then section 7 briefly reviews a few constructs such as multiscale, multiphysics, hybrid modeling approaches, hot spots and hot moments (HSHMs), and functional zonation that evolved over the years to transfer knowledge across sites. We then review legacy contamination and their coupled interactions with C, N, and S dynamics in the backdrop of the HSHM construct. After that, in section 8, we identify the current ecosystem science, eco-hydro-biogeochemistry, that links ecology, hydrology, and biogeochemistry. Following this, we highlight global water security issues and offer possible solutions in section 9. Finally, section 10 discusses the implications of scientific advances in ecosystem science and provides concluding thoughts and future research directions.

3. Description of the DOE's representative sites

To summarize the broad understanding that has emerged from DOE research, we choose seven DOE representative sites or testbeds: Savannah River Site in South Carolina, Oak Ridge Reservation in Tennessee, Hanford in Washington, Nevada National Security Site in Nevada, Riverton in Wyoming, and Rifle and East River in Colorado (figure 2). The Savannah River Site, Oak Ridge Reservation, Hanford, and Nevada National Security Site are DOE-EM sites. The Riverton and Rifle sites have been managed by the Office of Legacy Management (LM), which was established in 2003 to manage the remaining legacy of World War II (i.e. radioactive and chemical waste, contaminants, and hazardous material) at over 100 sites across the country. The East River is not associated with any DOE historical contamination, but represents evolution of DOE research toward broader watershed systems science. In addition, all of these sites were

later funded by the SBR program to study ecosystem function, thereby providing an excellent opportunity to learn how near-surface and subsurface legacy contaminant transport is modified by terrestrial and aquatic ecosystems.

The Savannah River Site, Oak Ridge Reservation, Hanford, Nevada National Security Site, and Riverton are respectively located in the South Atlantic-Gulf Region, Tennessee, Pacific Northwest, Lower Colorado regions, and Missouri regions. The Rifle and East River sites are located in the Upper Colorado River Basin, the principal water source in the southwestern USA. The Colorado River, one of the major rivers in the region, supplies water for 40 million people in seven states: Arizona, California, Colorado, Nevada, New Mexico, Utah, and Wyoming. These sites span across a wide range of hydro-climatic and geologic conditions. Table 1 describes contrasts in climate, geology, hydrogeology, surface water bodies, and groundwater depths across the representative sites.

Although several other DOE-funded testbeds have advanced the environmental sciences, they were not included in this review to keep the discussion manageable. This was particularly the case for sites with an extensive focus on aboveground processes, while this contribution's primary focus is to review the progress in understanding belowground processes and ecosystem function.

3.1. Savannah River Site

The Savannah River Site, located in south-central South Carolina, near Aiken, is an 800 square kilometer area where facilities were constructed in the early 1950s to produce special radioactive isotopes (e.g. Pu and tritium (^3H)) for the Department of Defense nuclear weapons stockpile. The major facilities constructed included production reactors, chemical processing plants, and solid and liquid waste storage sites. It is estimated that the Savannah River Site has approximately $172 \times 10^6 \text{ m}^3$ of groundwater, soil, and debris contaminated with metals, radionuclides, and organics (National Research Council 2000). Contamination of the environmental resources is the result of disposal practices conducted on-site. For example, the F-Area Seepage Basins (located in the north-central portion of Savannah River Site) consists of three unlined, earthen surface impoundments that received approximately 7.1 billion liters of acidic low-level radioactive (e.g. U, iodine (I)) waste solutions.

3.2. Oak Ridge Reservation Site

The Oak Ridge Reservation (formerly known as Clinton Engineer Works) was established in the early 1940s as part of the Manhattan Project on 239 square kilometers of land in eastern Tennessee, approximately 40 kilometers west of Knoxville, Tennessee. Its original missions, conducted at three large facilities, included U isotope enrichment, construction,

Table 1. Geologic and climatologic variability across the DOE sites (National Research Council 2000, Zachara *et al* 2013, Dam *et al* 2015, Dwivedi *et al* 2018b).

DOE site	Research focus	Climate	Geology and hydrogeology	Surface water body	Approximate depth to groundwater
Savannah River Site	Wetland hydro-biogeochemistry	Humid, subtropical; average annual rainfall 122 cm	Atlantic Coastal Plain with clay soils; the strata are deeply dissected by creeks, and most groundwater eventually seeps into and is diluted by creeks.	Savannah River and its tributaries	0 m–46 m
Oak Ridge Reservation	Stream corridor hydro-biogeochemistry	Humid, typical of the southern Appalachian region; average annual rainfall 138 cm	Valley and ridge province bordering the Cumberland Plateau; primary porosity is low, but fracture porosity present; high clay content; shallow water table.	Clinch River	1 m–37 m
Hanford Site	River corridor eco-hydro-biogeochemistry	Arid, cool, mild winters and warm summers; average rainfall 16 cm	Alluvial plain of bedded sediments with sands and gravels; Groundwater flows toward the Columbia River.	Columbia River	10 m–90 m
Nevada National Security Site	Hydro-biogeochemistry of actinides	Arid, mild winters and warm summers; average rainfall 13–32 cm	Alluvium, volcanic, and carbonate geology that is part of the Death Valley regional flow system.	Ephemeral streams, transient ponding, springs	210 m–610 m
Riverton Site	Floodplain hydro-biogeochemistry	Arid to semi-arid, steppe; average annual rainfall 25 cm	Gravel bed alluvial floodplain overlain by redox-active fine sediments; Wind River Formation.	Little Wind River	0 m–2.5 m
Rifle Site	Contaminants, C and N cycling	Semiarid; average annual rainfall 35 cm	Wasatch Formation overlain by Quaternary floodplain deposit.	Colorado River	1 m–5 m
East River	Watershed function	Arid to semiarid; annual rainfall 30 cm and 70–90 cm as snow	A diverse suite of Paleozoic and Mesozoic sedimentary rocks intruded by Tertiary igneous laccoliths and ore-rich stocks; Cretaceous Mancos Shale bedrock overlain by glacial moraine deposits.	East River and several tributaries representative of a headwater system	0 m–100 m

and operation of the world's first continuously operating nuclear reactor to demonstrate Pu production and separation, and radiochemical research and development. Today, the Oak Ridge Reservation covers an area of approximately 134 square kilometers and includes three major DOE complexes: the Oak Ridge National Laboratory (ORNL), the East Tennessee Technology Park, and the Y-12 National Security Complex (Y-12 NSC). Over the past 70 years, historical activities have resulted in significant releases of contaminants into the environment on the Oak Ridge Reservation, contaminating soils, sediments, groundwater, surface water, and biota. Waste disposal areas, unlined infiltration pits, trenches, and spills and leaks have created extensive subsurface contamination. Additionally, from 1950 through 1963, lithium (Li) isotope separation processes at the Y-12 NSC resulted in the release of about 212 000 kg of Hg into the environment, of which about 108 000 kg were estimated to have been lost to the East Fork Poplar Creek (EFPC). Both historical and ongoing Hg releases continue to negatively impact downstream waters, including many kilometers of the river–reservoir system outside the Oak Ridge Reservation.

3.3. Hanford Site

The Hanford Site, located in southeastern Washington State, was established by the federal government to conduct defense-related nuclear research, development, and weapons production activities. As a result of activities at the Pu production complex and processing facilities (chemical separations plants and solid–liquid disposal and waste storage sites), it is estimated that the 1517 square kilometer area has approximately $1400 \times 10^6 \text{ m}^3$ of groundwater, soil, and debris contaminated with metals, radionuclides, and organics (National Research Council 2000). Contamination of the sites' environmental resources occurred as a result of 1.3 trillion liters of wastewater from chemical processing operations being intentionally discharged into the ground through settling ponds and other subsurface drainage structures (DOE 1997). Additionally, DOE estimates that 67 of the 177 underground high-level radioactive waste storage tanks have leaked 3.8 million liters or more of the highly radioactive waste into the subsurface (DOE 1997). Most of Hanford's subsurface contamination is concentrated in two locations: the Columbia River Corridor (100 Area and 300 Area) and the Central Plateau (200 Area). The 544 square kilometer Columbia River Corridor contained the production reactors, several waste burial sites, and major research facilities (DOE 1996). The 194 square kilometer Central Plateau, near the middle of the Hanford Site, contained the chemical processing facilities for extracting U and Pu from irradiated reactor fuel, associated waste storage facilities (18 tank farms), and waste disposal facilities (i.e. surface settling basins and underground drainage cribs) (DOE 1996). It is estimated

that more than 220 square kilometers of groundwater at the Hanford Site is contaminated above current standards, mostly from operations in the 100 and 200 Areas. Section 5 briefly discusses the modeling of U contamination in the Hanford 300 Area and cesium (Cs) migration at the Hanford 200 Area.

3.4. Nevada National Security Site (formerly Nevada Test Site)

The Nevada National Security Site (NNSS), located in southern Nevada, was the primary site for USA's underground nuclear testing. It led to the deposition of substantial quantities of 43 different radionuclides into the environment (Smith *et al* 2003). While ^3H is the most abundant anthropogenic radionuclide deposited in the NNSS subsurface from an activity standpoint, Pu is the most abundant anthropogenic element by mass. Between 1951 and 1992, 828 underground nuclear tests (and 100 aboveground tests) were performed, and approximately 2.8 metric tons of Pu remains in the NNSS subsurface (DOE, Nevada Operations Office 2000). The site is approximately $\sim 145 \text{ km}$ northwest of Las Vegas, Nevada, and covers approximately 3500 square kilometers. The DOE-EM program has been developing a monitoring strategy that involves identifying contaminant boundaries, restricting access to contaminated groundwater, and implementing a long-term monitoring program.

3.5. Riverton site

The Riverton, Wyoming DOE legacy site is a riparian floodplain that was the location of a U and vanadium (V) ore processing mill that conducted operations between 1958 to 1963 (DOE-LM 1998). It is situated at $\sim 1500 \text{ m}$ elevation along the gravel bed of Little Wind River and has a semiarid to arid climate (25 cm mean annual precipitation). Temperatures exceed 0°C 154 days per year. The floodplain is vegetated with steppe flora, including sagebrush, grasses, and willows. Roots are relatively abundant, and robust upward solute transport through evapotranspiration has resulted in extensive evaporite mineralization of unsaturated soils (Dam *et al* 2015). The upper 2.5 m of soil is predominantly loam with abundant clay lenses, deposited abruptly over the underlying gravelly–cobble–sandy alluvial bed. Groundwater contains persistent U, molybdenum (Mo), and sulfate (SO_4^{2-}) contamination from an upgradient legacy U ore processing facility. Soils are loamy, but clay layers and sand lenses are present above and within the unconfined shallow sandy–gravelly aquifer (2.5 m–3 m below the ground surface). The site experiences intense seasonal redox activity triggered by rising and ebbing water tables and frequent flood events. Clay lenses exhibit molecular oxygen (O_2) depletion upon water saturation and the establishment of reducing conditions (including iron sulfide (FeS) precipitation). Significantly, reducing conditions extend into

the surrounding sandy aquifer and persist in proximity to clay lenses.

3.6. Rifle Site

The Rifle, Colorado DOE legacy site is a riparian floodplain that was the location of a former U and V ore processing facility that operated from 1924 through 1958 (DOE-LM 1998). The Rifle Site is approximately 750 m in length along the Colorado River shore and 250 m at the widest point. The former processing facility contained large piles of mill tailings on-site, from which residual U and other associated contaminants leached into the subsurface. All surface structures and contaminated soil were subsequently removed, and the site was capped with fill material and vegetated. However, the potential for infiltration of groundwater contaminants remained until that time (Williams *et al* 2011, Long *et al* 2012). Site-specific investigations revealed that the groundwater contained dissolved U at the micromolar concentration level. Other contaminants of concern were identified as As, selenium (Se), and V. Groundwater at the site flows through alluvial flood plain deposits, which are 6 m–7 m thick and underlain by the relatively impermeable Wasatch formation. Groundwater enters the alluvial aquifer from upgradient sources above the floodplain and exits into the Colorado River.

Given this background, earlier research was focused on *in situ* immobilization of contaminants. More recently, an increased focus was placed on constraining the exchanges and fates of different forms of C and N in river–floodplain settings because of their important roles in driving biogeochemical interactions with contaminants. More broadly, efforts were dedicated to developing a fundamental understanding of microbial communities and how they mediate biogeochemical cycles in the terrestrial subsurface. More than 15 years of focused investigations at the site have created a rich legacy of understanding subsurface contaminant transport and provided transferable insights.

3.7. East River Site

The East River Watershed, located in western Colorado, is a mountainous headwater testbed for exploring how climatic perturbations impact downgradient water availability and quality. The East River Watershed is located approximately 160 kilometers southeast of the Rifle Site and covers approximately 300 square kilometers. The East River is a pristine watershed. The Berkeley Lab's Watershed Function Scientific Focus Area (SFA; <https://watershed.lbl.gov/>) is examining the contribution to the riverine budgets of C and N that were neglected in previous assessments of contaminated sites like Rifle. The East River testbed began operation in 2014. Since then, it has enhanced process understanding of watershed dynamics and developed new approaches (such

as monitoring strategies using sensor networks and scale-adaptive approaches) to improve watershed-dynamics simulation (Hubbard *et al* 2018, Arora *et al* 2020).


4. Advances and innovations through DOE science in the last 20 years

DOE research has evolved in scale and complexity, beginning with fundamental microbiology and geochemistry, and then linking them into biogeochemistry. Further, this research expanded to hydro-biogeochemistry (figure 3) in unison with characterization and monitoring strategies. A detailed description of these is highlighted below.

4.1. Advances in subsurface microbiology

It is widely understood that microbial communities, not individual microbes, are the drivers of most subsurface biogeochemical cycles (figure 3(A)). Yet, in the early period of biological research at the Rifle Site, the focus was primarily on one family of bacteria, the *Geobacteraceae*, which have a laboratory-demonstrated capacity for U reduction (Gorby and Lovley 1992). This body of work motivated *in situ* acetate injection experiments to test the potential of a *Geobacter*-based bioremediation strategy at the Rifle Site (Anderson *et al* 2003). Proteomics of subsurface-derived cells (metaproteomics) were attempted to assay for *Geobacter*-specific proteins in amended aquifer groundwater. The only reference protein sequences available for this analysis were from *Geobacter* isolate genomes (Wilkins *et al* 2009). Thus, the effectiveness of protein identification was limited because the aquifer *Geobacter* genotypes differed from those in pure cultures.

The first proteomics experiment motivated one of the first cultivation-independent genomics (metagenomic) analyses of groundwater to improve the representation of *Geobacter* protein sequences. DNA from a *Geobacter* isolate not present at the Rifle Site provided an internal standard that was used to verify the accuracy of the recovered sequences. The experiment uncovered the very substantial microbial diversity of groundwater microbiomes. Draft genomes were reconstructed for over 80 bacteria, most of which were from previously unsampled or unknown lineages. Approximately half were from a group of novel bacteria whose gene content suggested they were symbionts of other microorganisms (Wrighton *et al* 2012). These groups were subsequently deeply sampled in a larger acetate injection experiment at the site and from unamended groundwater, and are now recognized as essentially ubiquitous in groundwaters worldwide (He *et al* 2021). The lineages share a common ancestry, so they were described as the Candidate Phyla Radiation (CPR) and were shown to comprise a substantial fraction (15% to ~50%) of all bacterial diversity (Brown *et al* 2015, Hug



et al 2016). The uniformly small genomes for CPR bacteria (Wrighton *et al* 2012, Kantor *et al* 2013) predicted their small cell size and thus potential to pass through the filters routinely used for cell recovery. Imaging of cells from the smallest filtrate fraction from acetate-amended groundwater revealed that the cells are around the theoretical minimum size for life (Luef *et al* 2015).

Also first detected in Rifle groundwater were novel archaea with genomic features reminiscent of CPR bacteria. These diverse archaea, now part of the DPANN group, substantially extended the breadth of known archaeal diversity (Castelle *et al* 2013). Given that both CPR bacteria and DPANN archaea are generally predicted to grow as surface-attached symbionts of other cells, it has been inferred that they have the potential to impact biogeochemical cycles through their impacts on host cells as well as via their own capacities for C, H, N, and S compound transformations. Subsequent research identified numerous bacteriophages (phages), the virus of bacteria, in subsurface communities (Al-Shayeb *et al* 2020), bringing them into focus as important contributors to C compound turnover.

Sampling of subsurface microbial communities from pumped pore fluids overlooks particle-associated cells. This motivated the study of sediment-associated consortia and uncovered very significant microbial diversity (Castelle *et al* 2013). One of the first applications of long-read sequencing to complex microbiomes revealed the presence of thousands of different organisms, most of which are at similar, very low abundance levels in Rifle sediments (Sharon *et al* 2015). Metagenomic analyses of saturated whole sediment samples recovered from 4, 5, and 6 m depths identified numerous new lineages,

one of which is the bacterial phylum Zixibacteria. From a complete, curated genome, Zixibacteria were predicted to have an incredible diversity of metabolic capacities that enable growth despite shifts in conditions and the available aquifer resources (Castelle *et al* 2013). Another study targeting sediment-associated Chloroflexi revealed, among other traits, an unexpected pathway for carbon dioxide (CO₂) fixation that brings into focus the potential for this process in C cycling in the aquifer (Hug *et al* 2013).

A study that utilized groundwater and sediment metagenomes sampled from below and above the water table indicated that microbial cohorts consistently establish across the aquifer (Hug *et al* 2015). This, along with the desire to develop models for integrated microbial community function (Zhuang *et al* 2011), motivated a major genomes recovery effort. Genomes from over 2000 different microbial community members were analyzed simultaneously (an unprecedented effort). The major conclusion from this study was that biogeochemical cycles are attributed not to single organisms with capacities for specific pathways but to consortia whose metabolisms are interlinked by handoffs of pathway intermediates (Anantharaman *et al* 2016). This ecosystem structure probably confers resilience and has since been documented by genomics-based studies in other ecosystems, including soil (Diamond *et al* 2019).

Important limitations of the Rifle research site were its comparatively small size, lack of access to undisturbed soil and riparian zone sediments, and limited vegetation, slope, and aspect types. Thus, methods developed at the Rifle Site were scaled up to tackle the challenge of watershed ecosystem-scale microbial biogeochemical process analyses that incorporate genome-derived information. Initial studies

focused on the soils surrounding the meanders of the East River Site in Colorado. Motivated by the need to link micron-scale microbial analyses to a transect that is many kilometers in length, it was hypothesized that soils within the arcs of successive meanders would represent a system representative of meanders along the river length.

Metagenomic datasets were generated from a sampling grid at three sites in the river's upper, middle, and lower reaches. Extensive genome recovery enabled documentation of biogeochemically important capacities across each site. A core floodplain microbiome was identified and found to be enriched in capacities for aerobic respiration, aerobic carbon monoxide oxidation, and thiosulfate oxidation. Metatranscriptomic data revealed that the most highly transcribed genes were *amoCAB* and *nxrAB* (for nitrification). Low soil organic C correlates with the high activity of genes involved in methanol, formate, sulfide, hydrogen, and ammonia oxidation, nitrite oxidoreduction, and nitrate and nitrite reduction. Overall, it was concluded that the meander-bound regions serve as scaling motifs that predict aggregate capacities for biogeochemical transformations in floodplain soils (Carnevali et al 2021). Similar research has addressed patterns of organism distribution and function across hillslopes and under four different vegetation types.

4.2. Advances in geochemistry and biogeochemistry

DOE's research in water and energy security arose from its historical and present-day need to resolve the many environmental management issues across its vast nuclear facilities, and its responsibility to develop a long-term solution to the USA's growing nuclear waste stockpile associated primarily with civilian nuclear energy production. The range of radiologic contaminants quite literally spans the entire periodic table: from ^3H to the short-lived heavy elements (e.g. curium-244 (^{244}Cm)) (Kurosaki et al 2014). More traditional contaminants of concern include Cr, Hg (discussed in section 6), and highly radioactive Cs and radioactive U (discussed in section 5.2), and more recently C and nutrients, including N and phosphorus (P). Below, we briefly describe DOE advancements in understanding the biogeochemical processes impacting contaminant fate and transport, as well as the major elements and nutrients (i.e. C, N, Fe, S, and manganese (Mn)) that drive subsurface biogeochemistry.

4.2.1. Contaminant biogeochemistry in redox-dynamic environments

U is a major risk driver at many of DOE's legacy waste sites, and DOE-funded programs have made major contributions to our knowledge of molecular species and processes controlling the movement of U in natural waters. The mobility of U in the DOE legacy

waste sites is largely controlled by its speciation. The speciation includes dissolved, adsorbed, and mineralized species, predominantly in the +4 or +6 valence states. U^{V} is generally unstable and disproportionates to U^{VI} and U^{IV} (O'Loughlin et al 2011); however, U^{V} has been detected in environmentally relevant systems (Ilton et al 2005, Nico et al 2009b), and stable U^{V} associated with Fe^{II} -containing clays has been reported (Boyanov et al 2016, 2017a). Hexavalent U is stable and soluble under oxic conditions as the uranyl cation (UO_2^{2+}) (Ragnarsdottir et al 2000). Tetravalent U is stable and sparingly soluble under reducing conditions, but can be mobilized as complexes with organic or inorganic C (Frazier et al 2005, Luo and Gu 2009, Stoliker et al 2013).

U^{VI} can be reduced by microbial organisms (Wall and Krumholz 2006, and references therein) and structural Fe^{II} in minerals (O'Loughlin et al 2003a, 2010, Ilton et al 2005, Ithurbide et al 2009, Veeramani et al 2011, Singer et al 2012, Latta et al 2012b, Boyanov et al 2017a). These reactions generally produce amorphous U^{IV} products (Kelly et al 2008, Sharp et al 2011, Latta et al 2012a, Bargar et al 2013, Wang et al 2014a, Morin et al 2016, Bone et al 2017a, 2017b, 2020, Stetten et al 2018). For example, adsorbed U^{IV} was found to be the dominant form of U in naturally reduced sediments at legacy DOE sites across the Upper Colorado River Basin (Noël et al 2017a, 2017b). NOM functional groups, phosphate or phosphoryl groups, and high-affinity titanium (Ti)- or Fe-based binding sites in minerals have been cited as important ligands for U^{IV} in natural systems (Fletcher et al 2010, Boyanov et al 2011, 2017b, Sivaswamy et al 2011, Veeramani et al 2011, Stylo et al 2013, Latta et al 2014, Wang et al 2015, Bone et al 2017b, 2020). U redox cycling also can lead to the production of U^{V} in Fe^{III} oxyhydroxides (Kerisit et al 2011, Massey et al 2014, Dewey et al 2020). The finding that sorbed species are a major mode of occurrence of U^{IV} was initially unexpected and has subsequently informed the development of reactive transport models (Yabusaki et al 2017).

Oxidation of U^{IV} by O_2 has been studied extensively for uraninite (Gu et al 2005, Boyanov et al 2007, Senko et al 2007, Ulrich et al 2009, Lezama-Pacheco et al 2015). Oxidation of UO_2 by Fe^{III} oxides and clays has also been documented within the DOE programs (Sani et al 2005, Ginder-Vogel et al 2006, Senko et al 2007, Stewart et al 2013). Oxidation of non-uraninite U^{IV} species by O_2 has also been investigated to a limited extent, producing mixed results regarding their stability relative to uraninite (Sharp et al 2011, Cerrato et al 2013, Bi et al 2016, Latta et al 2016). Another recent finding is that the presence of reduced C and Fe phases inhibits oxidative release of non-uraninite U^{IV} (Newsome et al 2015, Bone et al 2017a).

Unlike U, which is regulated at picomolar concentrations, many radionuclides are regulated at femtomolar concentrations (e.g. The United

States Environmental Protection Agency (EPA) Maximum Contaminant Levels for drinking water of 6.4×10^{-16} , 1.7×10^{-14} , and 3.7×10^{-15} mol L⁻¹ for strontium-90 (⁹⁰Sr), ¹³⁷Cs, and ²³⁸Pu, respectively) (Deblonde *et al* 2020). The body of environmental radiochemistry research supported by the DOE is much too vast to summarize here; however, we have selected a few examples to highlight key advances in this area. Much of DOE's research in environmental radiochemistry has focused on a process-level understanding of radionuclide reactive transport. Examination of colloid-facilitated Pu transport revealed the importance of both organic (Xu *et al* 2008, Zhao *et al* 2011) and inorganic colloids (Zhao *et al* 2020), the nature of these colloid associations (Powell *et al* 2011), the importance of redox cycling (Pan *et al* 2021), and the sorption-desorption kinetics that limit colloid-facilitated transport (Begg *et al* 2017, 2018). A combination of molecular dynamics simulations revealed the importance of Cs interaction with minerals via ion exchange (Zaunbrecher *et al* 2015) along with experiments that determined the retardation properties controlling its migration (Zachara *et al* 2002, Steefel *et al* 2003, 2005, Durrant *et al* 2018). The complex chemistry and reactive transport of I were found to be controlled by a combination of redox chemistry and organic matter associations (Neeway *et al* 2019). Indeed, the importance of redox conditions in the transport behavior of radionuclides is a theme that spans across several radionuclides of concern to the DOE (¹²⁹I, technetium (Tc), U, neptunium (Np), and Pu) (O'Loughlin *et al* 2011). Section 5.2 describes the reactive migration of U and Cs at the Hanford and Rifle sites further illustrating the importance of redox conditions and demonstrating the comprehensive and rigorous treatment of geochemistry in contamination modeling, as part of the simulation capability development.

In addition to U and other radionuclides, Cr is a contaminant of concern at many DOE legacy waste sites. Cr is a naturally occurring, redox-active transition metal that can exist in a range of oxidation states, with Cr^{III} and Cr^{VI} being the most stable in environmental systems. Furthermore, Cr^{VI} is one of the most mobile forms of Cr, which is transported in surface and subsurface waters as a negatively charged oxyanion. Although hexavalent Cr is generally more soluble, mobile, and toxic, Cr^{III} is significantly less soluble than Cr^{VI} and is an essential micronutrient. At DOE sites, such as the Hanford 100 Area (near reactor D, see www.hanford.gov/page.cfm/100Area), sodium dichromate was used during reactor operations to retard corrosion in reactor cooling systems. Consequently, Cr was released to the environment by spills and/or leaks from pipes, resulting in Cr^{VI} contamination in the vadose zone and groundwater adjacent to the Columbia River. For over two decades, DOE has performed groundbreaking research on factors that control Cr fate and transport

at the Hanford 100 Area. For example, researchers investigated Cr interactions with sediments (Zachara *et al* 1987, 1988, Beller *et al* 2014) and the use of various forms of Fe and biostimulation to convert Cr^{VI} to Cr^{III} (Powell *et al* 1995, Fendorf and Li 1996, Jardine *et al* 1999, Mayes *et al* 2000, Hazen and Tabak 2005), illustrating techniques for transforming mobile Cr^{VI} to immobile Cr^{VI} in subsurface systems.

The fate and mobility of As in a heterogeneous redox environment are difficult to predict because it is influenced by mineralogy, chemical speciation, and biological processes. Arsenic has five oxidation states (−III, −I, 0, III, V), with As^{III} and As^V being the most prevalent oxidation states in surface and groundwater, where it typically is present as oxyanion (arsenate and arsenite) and thioanion species. Thioarsenic can dominate in sulfidic environments that contain large amounts of reduced S, such as organic-rich surface water and groundwater (Stucker *et al* 2014, Boye *et al* 2017, Kumar *et al* 2020a). Stucker *et al* (2013, 2014) observed the presence of thioarsenic species under SO₄^{2−}-reducing conditions while conducting a biostimulation experiment at the Rifle Site in Colorado to treat the U-contaminated groundwater plume. NOM is also an important consideration in understanding As fate and mobility in environmental systems, such as wetlands, riparian areas, and groundwater aquifers, where redox conditions vary over space and time. This type of redox heterogeneity can have an outsized influence on As speciation, release, and retention. Recently, Kumar *et al* (2020a) showed that As release and retention in natural aquifer sand is governed by the presence of thin organically rich clay lenses, which generate redox heterogeneities in sediments and thus alter the relative concentrations of Fe and S and promote the formation of mobile thioarsenate species.

4.2.2. Biogeochemical critical elements that mediate contaminant behavior

Mn is one of the most abundant trace metals in soils, sediments, and natural waters. Its complex redox chemistry, consisting of three commonly occurring oxidation states of Mn^{II}, Mn^{III}, and Mn^{IV}. The Mn^{II} state is highly soluble, while the Mn^{III} and Mn^{IV} states are sparingly soluble and occur most frequently as mixed hydr(oxides). While oxidation of Mn^{II} by O₂ is highly thermodynamically favorable, it is kinetically limited, leading to significant concentrations of dissolved Mn in many natural waters and leading control of Mn oxidation by a number of key processes. The most dominant of these is biological Mn oxidation, in which microorganisms including both bacterial and fungi catalyze the oxidation of Mn^{II} to Mn^{III}/Mn^{IV} (Spiro *et al* 2010, Santelli *et al* 2011, Geszvain *et al* 2012). These organisms oxidize Mn directly through enzymes such as multicopper oxidase as well as indirectly through the production of reactive oxygen species (ROS) (Hansel *et al* 2012). ROS

produced abiotically, e.g. through photochemistry or Fenton chemistry, also provide an abiotic pathway to Mn^{II} oxidation (Nico *et al* 2002, Learman *et al* 2011, van Genuchten and Peña 2016). The strong coupling of the Mn cycle with the ROS cycle is also thought to protect organisms from ROS-driven oxidative damage (Spiro *et al* 2010).

The oxidation of Mn^{II} can produce a wide variety of Mn (hydr)oxides with different structures and reactivity (Bargar *et al* 2009, Tan *et al* 2010, Droz *et al* 2015, Ling *et al* 2020). Given the ubiquitous nature of Mn (hydr)oxides, they play important roles at DOE sites. Some of the most common Mn (hydr)oxides have layered type structures consisting predominantly of MnO_6 octahedra, but also contain multiple vacancies and defects, including inclusions of Mn^{III} centers (Ilton *et al* 2005, Bargar *et al* 2009, Spiro *et al* 2010). The domain particle size is frequently on the nanoscale, making them natural nanoparticulate materials (Bargar *et al* 2009). Overall, Mn (hydr)oxides are both powerful oxidants of other elements and strong, high surface area adsorbents. In terms of sorption behavior, they have been shown to strongly sorb other key elements, including cobalt (Co), zinc (Zn), nickel (Ni), Mn, copper (Cu), Cr, cadmium (Cd), lead (Pb), and U (Duckworth *et al* 2009, Kwon *et al* 2009, 2013, Wang *et al* 2013a, Fuller and Bargar 2014, Peña *et al* 2015, Simanova *et al* 2015, van Genuchten and Peña 2016). This sorption behavior is frequently associated with vacancy sites within the oxides and can also include sorption at edges and structural incorporation. Mn (hydr)oxides frequently act as oxidants of other environmentally important metals and metalloids, including As, Cr, S, U, Tc, Co (Nico *et al* 2009a, Plathe *et al* 2013, Wang *et al* 2013a, 2013b, 2014a, 2014b, Duckworth *et al* 2014, Fan *et al* 2014, Tang *et al* 2014, McClain *et al* 2017). The oxidation of these species by Mn (hydr)oxides is important in the efficacy of contamination remediation approaches focused on reductive immobilization of critical elements. In the case of Cr, Mn (hydr)oxides appear to be the critical pathway by which Cr^{VI} is generated in natural systems (Plathe *et al* 2013, Wang *et al* 2013b, 2014b, Keiluweit *et al* 2015, Hausladen and Fendorf 2017, McClain *et al* 2017). More recently, Mn has been shown to play an important role in natural C cycling both in waters as well as soils and sediments, whereby either through direct reaction with Mn (hydr)oxides or by biological utilization of an Mn complex, organic material is oxidatively decomposed into smaller molecules (Keiluweit *et al* 2015, Estes *et al* 2017, Jones *et al* 2018, 2020). Microorganisms can directly utilize Mn (hydr)oxides as terminal electron acceptors, thereby coupled with C and other elemental redox cycles.

Fe oxides and Fe-bearing clay minerals are common constituents of soils and sediments at DOE sites. The biogeochemistry of Fe in most aquatic and terrestrial environments is driven largely by microbial

activity, particularly in Fe-rich soils and sediments where Fe redox cycling by microorganisms is a significant component of C cycling and energy flux (Nealson and Saffarini 1994, Roden and Wetzel 1996, Lovley 2000). As a group, dissimilatory iron-reducing bacteria (DIRB) can use a wide range of Fe^{III} forms as terminal electron acceptors for anaerobic respiration, including soluble Fe^{III} complexes, Fe^{III} oxides, and clay minerals containing varying amounts of structural Fe^{III} (Fredrickson *et al* 1998, Zachara *et al* 1998, Kukkadapu *et al* 2001, O'Loughlin *et al* 2021). DIRB activity can yield a suite of Fe^{II} species including soluble Fe^{II} complexes, Fe^{II} complexes with the surfaces of organic and inorganic solid phases, and a host of mineral phases containing structural Fe^{II} (Lovley *et al* 1987, O'Loughlin *et al* 2013, 2019, Dong *et al* 2020). Many of these Fe^{II} species effectively reduce a wide range of organic and inorganic contaminants of concern at DOE sites, including chlorinated hydrocarbons, nitrate, Cr^{VI} , U^{VI} , Tc^{VII} , Np^{V} , and Pu^{V} (Kelly *et al* 2002, Bond and Fendorf 2003, O'Loughlin *et al* 2003a, 2003b, 2020, Fredrickson *et al* 2004, Kemner *et al* 2004, Peretyazhko *et al* 2008, Wiatrowski *et al* 2009, Felmy *et al* 2011, Veeramani *et al* 2011, Latta *et al* 2012a). Finally, Fe^{II} (bio)oxidation can result in the formation of a variety of Fe^{III} -bearing minerals, depending on geochemical conditions.

Because of the relative insolubility of most Fe-bearing minerals in typical aquatic and terrestrial environments (i.e. circumneutral pH), their use for respiration by DIRB and iron-oxidizing bacteria (IOB) as terminal electron acceptors and donors, respectively, requires different mechanisms for electron transfer relative to soluble terminal electron acceptors/donors that are easily transported into the cell (e.g. O_2 , NO_x^- , SO_4^{2-}). DOE-funded research has advanced understanding of extracellular electron transfer to solid-phase electron acceptors. Some DIRB such as *Geobacter* and *Shewanella* can transfer electrons directly to Fe^{III} oxide surfaces by means of reductases located on their outer cell membrane (Shi *et al* 2009) or via electrically conductive pili or nanowires (Reguera *et al* 2005, Gorby *et al* 2006). The need for physical contact between Fe^{III} oxide minerals and microbial cells, however, can be readily overcome. The dissolution of Fe^{III} oxides is promoted by exogenous and endogenous ligands, and the resulting soluble Fe^{III} complexes can diffuse away and be reduced by DIRB at a distance (Nevin and Lovley 2002, Taillefert *et al* 2007). Likewise, the transfer of electrons from the cell to external electron acceptors (e.g. Fe^{III} oxides) can be facilitated by soluble electron shuttles, i.e. compounds that can be reversibly oxidized and reduced, which include a wide variety of endogenous and exogenous organic and inorganic compounds, including quinones, flavins, humic substances, and reduced S species (Nevin and Lovley 2000, 2002, Royer *et al* 2002, O'Loughlin 2008, Roden *et al* 2010, Flynn *et al* 2014).

Although S is often less abundant than Fe in soils and sediments at DOE sites, its biogeochemical transformations are more complex because of the variety of S oxidation states (VI, V, IV, III, II, I, 0, $-I$, and $-II$). In oxic environments S^{VI} (e.g. SO_4^{2-}) is typically the most thermodynamically stable valence state, while S^{-II} (e.g. H_2S and metal sulfides) is the most stable in anoxic environments. Unlike Fe^{III} , S^{VI} as SO_4^{2-} is a highly soluble anion found commonly in groundwater, as a consequence of the dissolution of SO_4^{2-} minerals or as a product of the oxidation of iron sulfide minerals (e.g. pyrite). Like DIRB, dissimilatory sulfate-reducing bacteria (DSRB) and archaea are anaerobes that can obtain energy by coupling the oxidation of organic compounds or molecular H_2 with the reduction of SO_4^{2-} to sulfide. This process is a key component of S and C cycles in anoxic soils and sediments (Meronigal *et al* 2004). In addition, sulfide generated by DSRB can reduce Fe^{III} oxides (Kwon *et al* 2014a, 2014b, Johnson *et al* 2021), resulting in the oxidation of sulfide, primarily to elemental S (as molecular S^0 and polysulfides) with minor amounts of thiosulfate. These pathways are important links coupling Fe and S biogeochemical cycles (Howarth *et al* 1992, Lovley 1993, Nealson and Saffarini 1994, Flynn *et al* 2014, Hansel *et al* 2015).

4.2.3. C and N biogeochemistry

C biogeochemistry in soils is linked with redox reactions (e.g. aerobic respiration, denitrification, methanogenesis) in a simplified manner by conceptualization of a single solid phase soil organic matter (SOM) that provides a sustained supply of dissolved organic carbon (DOC). However, DOC availability that drives redox reactions is not unlimited and is also influenced by SOM dynamics. The traditional modeling approach for belowground C cycling does not explicitly consider the primary underlying processes and agents (e.g. microbes, aggregation) important for SOM cycling. Instead, SOM is classified into multiple C pools based on their degradability and is qualitatively named—for instance, recalcitrant versus labile or active pools. An emergent modeling approach recognizes that SOM comprises many complex molecules (Schmidt *et al* 2011, Dwivedi *et al* 2019) and that SOM decomposition is a function of a wide range of ecosystem properties and mechanisms such as temperature, thermodynamics, redox status, moisture content, exoenzyme production, decomposition, organo-mineral interactions, microbial necromass, ecology of belowground biota, and root contributions, as well as the mobilization of nutrients (as shown in figure 3(C), Schmidt *et al* 2011, Keiluweit *et al* 2016, Boye *et al* 2017, Dwivedi *et al* 2019, and references therein). As a result, the DOC supply that drives redox reactions varies significantly in space and time, and has implications for redox processes.

Motivated by this emerging understanding, Riley *et al* (2014) developed a complex reaction network

of SOM, including above- and belowground organic inputs, multiple DOC species, and heterotrophic microbes. They captured vertical SOM variability reasonably well. However, they did not include explicit organo-mineral interactions in their reaction network, and used a Langmuir isotherm to represent sorption (Riley *et al* 2014), which does not adequately account for the suite of minerals present in the subsurface. Dwivedi *et al* (2017a) later enhanced this reaction network by including a Surface Complexation Model (SCM) and demonstrated controls exerted by plant roots and mineral surface area on availability and persistence of DOC in the soil.

Maggi *et al* (2008) developed a reaction network primarily consisting of ammonia oxidation and denitrification to examine N cycling in near-surface soil. Maggi *et al* (2008) did not consider other redox species, like Fe and S, in their reaction network as their investigation was limited to N cycling for an agricultural field. However, Arora *et al* (2016b) developed a reaction network representing a redox staircase for the Rifle Site, where redox species (e.g. Fe, S) and naturally reduced sediments were predominantly present. Dwivedi *et al* (2018a) subsequently used this reaction network at the Rifle Site and investigated the N dynamics in a floodplain environment. They demonstrated that, although the Rifle Site experiences little precipitation (being in an arid climate), short and episodic summer rain events led to the formation of HSHMs of N species due to nitrification resulting from the percolation of oxic water. Yabusaki *et al* (2017) also examined N biogeochemistry at the Rifle Site; however, they did not explicitly consider nitrification processes and instead focused on the reductive pathways. Moving away from a contaminated site, Dwivedi *et al* (2018b) also examined N biogeochemistry within intra-meander regions of the East River site. The East River, a pristine watershed, is low in nitrate. N cycling in the East River is primarily linked with snowmelt, vegetation, and atmospheric deposition (Newcomer *et al* 2021).

4.2.4. Coupling of contaminant and nutrient redox cycles

Redox-active U in natural sediments is coupled to the C cycle through metabolic oxidation of organic matter by microbes to CO_2 . Electrons released from this process are transferred to U^{VI} , reducing it to U^{IV} (see section 4.2.1). This process is the basis for stimulated bio-reduction, in which exogenous DOC is injected into groundwater, stimulating the reduction of U^{IV} and resulting in net attenuation of dissolved U (Williams *et al* 2011, Long *et al* 2012). Another mode by which contaminant and nutrient cycles interact is via complexation reactions with functional groups produced via metabolic and anabolic processes. For example, NOM coatings, abundant on clay and other minerals, strongly complex U^{IV} (see section 4.2.1). As pointed out elsewhere in this paper, methyl groups

are profoundly important as complexing ligands Hg in natural waters, and organic functional groups containing reduced S are important as complexing agents for As (see section 4.2.1) and Hg (see section 6).

Nitrate-facilitated oxidation of U^{VI} has been extensively investigated and provides numerous examples of strong coupling between the U and N cycles, as well as of the impacts of the Fe and S cycles on U-N redox coupling. Researchers have noted that when groundwater enriched in nitrate is amended with DOC (e.g. acetate), U^{VI} will not be reduced until nitrate has been consumed, and metal reduction or SO_4^{2-} reduction commences (Finneran *et al* 2002, Senko *et al* 2002, Istok *et al* 2004). Furthermore, U^{VI} can be recovered from aquifer sediments that have previously been reduced when groundwater is amended with nitrate (Finneran *et al* 2002, Istok *et al* 2004, Wu *et al* 2010). There are multiple pathways through which nitrate reduction can lead to U^{IV} oxidation. It has been suggested that some bacteria can couple denitrification to U^{IV} oxidation (Finneran *et al* 2002, Beller 2005). However, nitrate-dependent microbial oxidation of U^{IV} may be slower than abiotic oxidation of U^{IV} under nitrate-reducing conditions (Senko *et al* 2005b); for instance, U^{IV} is oxidized more quickly by nitrite formed during denitrification (Senko *et al* 2002). U^{IV} is oxidized most rapidly by Fe^{III} oxides that are produced by the oxidation of Fe^{II} by nitrite (Senko *et al* 2005a). The extent (Ginder-Vogel *et al* 2006) of U^{IV} oxidation by Fe^{III} (hydr)oxides depends strongly on the type of Fe^{III} (hydr)oxide; nanocrystalline/amorphous Fe^{III} oxides may react rapidly with U^{IV} due to high surface areas (Senko *et al* 2005a). In support of these conclusions, recent studies performed under diffusion-limited conditions suggest that denitrification occurs rapidly in soils and aquifers that host reduced U and is unlikely to overwhelm the reducing capacity of sediments that resides in the form of Fe^{II} , sulfide, and other reduced species.

Although O_2 can oxidize U^{IV} , nitrate-dependent U^{IV} oxidation has been shown to occur much more rapidly (Moon *et al* 2007, 2009). Using sediments from the background area of the Rifle Site, Moon *et al* (2007, 2009) demonstrated that slower oxidation of U^{IV} under oxic conditions was due to O_2 reacting quickly with numerous other reduced sediments species. In contrast, nitrate was less reactive towards the reduced species and could travel farther into the sediments and reach a greater total amount of U^{IV} . In fact, O_2 breakthrough did not occur for several weeks, whereas nitrate breakthrough occurred within hours (Moon *et al* 2007, 2009). Furthermore, in the presence of FeS, only 1% of U in the sediment columns was solubilized in the presence of O_2 over the same time period as 11%–60% U was solubilized in the presence of nitrate (Moon *et al* 2009). Buffering of U^{IV} against oxidation by reduced species including Fe^{II} , S^{II} , and organic C has been suggested by

several other researchers (e.g. Abdelouas *et al* 1999, N'Guessan *et al* 2010, Spycher *et al* 2011).

4.3. Advances in hydro-biogeochemistry

Hydro-biogeochemistry is essential for tracking and monitoring contaminants and nutrients that affect river water quality through subsurface geochemical fluxes. The subsurface geochemical behavior of contaminants and nutrients differs across sites due to distinct environmental, geological, and hydroclimatic conditions (Zachara *et al* 2013, Neeway *et al* 2019). In particular, sediments, mineral composition, geology, and hydroclimatic conditions give rise to contrasting mechanisms controlling plume persistence in the subsurface and riparian corridors. For example, several studies have demonstrated that the effect of mineral surfaces and sediments is important for the transformation and export of geochemical species (Keppler *et al* 2000, Fox *et al* 2012, Allard and Gallard 2013, Maher *et al* 2013, Cumberland *et al* 2016, Dwivedi *et al* 2017a, 2019), while hydrology plays a critical role in mobilizing contaminants (Li *et al* 2010, Flores Orozco *et al* 2011, Williams *et al* 2013, Dangelmayr *et al* 2017, Jemison *et al* 2020).

In a comparative study, Zachara *et al* (2013) outlined key differences in the geochemical nature of residual contaminant U at the Rifle and Hanford 300 Area sites. They suggested that the major differences between the sites and the kinetic processes influencing the U distribution in space primarily resulted from the spatial geochemical heterogeneity of detrital organic matter and the magnitude of groundwater hydrologic dynamics controlled by river stage fluctuations.

At DOE sites, including the Rifle and Columbia River Corridor at Hanford, it was found that the interplay between groundwater dynamics and river stage oscillations may result in gradient reversal, changes in flow velocities, redox zonation, and alterations in groundwater composition (Greskowiak *et al* 2010, Yabusaki *et al* 2011, Lezama-Pacheco *et al* 2015, Dwivedi *et al* 2018a, Noël *et al* 2019). The relative variations in groundwater levels and river stages can alter the mixing zone's extent, where groundwater meets the surface waters and exchanges solutes (Ellis *et al* 2007, Fritz and Arntzen 2007). The mixing zone may span over distances of the meter- to kilometer scales (Dwivedi *et al* 2017b, 2018b, Zachara *et al* 2020). For example, Arora *et al* (2016b), Dwivedi *et al* (2018a), and Yabusaki *et al* (2017) demonstrated that the subsurface geochemical exports, and thus river water quality, were influenced by the spatial geochemical heterogeneity of detrital organic and temporal variability in groundwater flow direction at the Rifle Site. Similarly, the Columbia River Corridor site at Hanford also experiences frequent hydrologic gradient reversals, primarily due to dam operations and distinct subsurface hydrogeology control (Shuai *et al* 2019). Overall, the gradient reversal at sites like

Rifle and Hanford can result in episodic oxic and anoxic conditions in the subsurface (Yabusaki *et al* 2011, Lezama-Pacheco *et al* 2015, Dwivedi *et al* 2018a, Chen *et al* 2021). Consequently, changes in the oxic environment due to groundwater mixing and microbially mediated sequential biogeochemical processes can lead to distinct redox gradients through a succession of electron acceptors.

4.3.1. Influence of hyporheic exchange flows and water-table fluctuations on mobilizing contaminants

The hyporheic zone, an active dynamic ecotone where nutrient-rich groundwater and oxic river water interact, extends below the saturated sediments adjacent to the flowing channels of streams and rivers and signifies the elevated reaction rates for nutrients and contaminants (e.g. Mulholland *et al* 2008, Dwivedi *et al* 2018b). The hyporheic exchange flows, referring to the water movement between hyporheic zones and river channels, are important controls on contaminant and nutrient transformations and ultimately export to downstream surface water bodies. To illustrate this further, the dynamic redox behavior in the hyporheic zone influences the fate and transport of Cr, a common contaminant in subsurface and surface water. Liu *et al* (2017) investigated Cr behavior in the Columbia River hyporheic zone at the Hanford Site. They found that the hyporheic zone acts as a natural redox barrier for reductively immobilizing Cr under dynamic hydrological conditions.

In the Upper Colorado River Basin, seasonal water-table elevations can vary by up to 2 m over the course of the year (Noël *et al* 2019, Tolar *et al* 2020). During wet spring conditions, invasion of water into sediment pore space displaces air (and O₂), allowing reducing conditions to develop in shallow sediments that can persist into late summer in fine-grained clay-rich sediments (Tolar *et al* 2020). Subsequently, drainage of water from sediments in late summer allows air to perfuse in, facilitating re-oxidation of U^{IV}, as well as Fe^{II} and sulfides (Lezama-Pacheco *et al* 2015, Lefebvre *et al* 2019, Noël *et al* 2019). Under repeated annual cycles, complex patterns of U, as well as S and Fe redox cycles, overlap, creating distinctive signatures of redox conditions (Noël *et al* 2017a, 2017b).

4.3.2. Influence of hydro-climatic conditions on contaminant and nutrient dynamics

Hydro-climatic conditions specific to precipitation and evapotranspiration are equally important from the contaminant and nutrient dynamics standpoint. Frequent and intense rainfall in subtropical and humid climates like the Savannah River Site leads to prolonged flooding and stormwater infiltration (O'Reilly *et al* 2012). Changing hydro-climatic and geochemical conditions and wet-dry redox cycles exert significant controls on contaminant cycling and mobilization-immobilization in the subsurface (Noël

et al 2017b, Song *et al* 2018). For example, redox processes such as nitrification can be significant for mobilizing contaminants (e.g. U), particularly when oxic river water or rainwater infiltration alters the redox gradients (Bone *et al* 2017a, Dwivedi *et al* 2018a, Cardarelli *et al* 2020, Tolar *et al* 2020). Noël *et al* (2019) showed that U accumulated as U^{IV} in shallow riparian floodplain sediments at the contaminated DOE Shiprock Disposal Site in New Mexico under water-saturated spring conditions, but was oxidized and precipitated as U^{VI} carbonates or silicates during summer drought.

Climate change is causing precipitation to decrease and evapotranspiration to increase in the Upper Colorado River Basin (Christensen and Lettenmaier 2007, Prein *et al* 2016), leading to decreasing water tables and exposure of reduced sulfidic sediments to oxygenated conditions. It is thus plausible that climate change will drive regional remobilization of accumulated U and other metals. In support of this conclusion, Janot *et al* (2016) estimated that oxidative release of U from a single moderate-sized naturally reduced zone at the Rifle Site could sustain a U plume at the regulatory concentration limit (0.044 mgU L⁻¹) for hundreds if not thousands of years.

Developing insights into the interplay of hydrology, geochemistry, and biology requires developing insights into processes controlling the cycling of contaminants, nutrients, and elements such as oxygen (O), Fe, S, C, and N in the groundwater and floodplain. High-fidelity multicomponent reactive transport modeling has been found to advance a robust predictive understanding of this interplay across the DOE representative sites (Yabusaki *et al* 2011, Li *et al* 2017, Dwivedi *et al* 2018a, Steefel 2019). Significant progress has been made in modeling complex multidirectional flow processes and their biogeochemistry interactions using these DOE representative sites. We describe these modeling developments in greater detail in section 5.

4.4. Characterization and monitoring strategies

Over the last several decades, there have been significant advances in the characterization of above- and belowground properties and dynamics necessary to parameterize, calibrate, and validate reactive transport models. At the point and single-borehole scales, there have been rapid advances in *in situ* sensors to measure groundwater and surface qualities, such as soil moisture, pH, water table, and electrical conductivity. These sensors are typically placed inside wells to continuously monitor groundwater or surface water.

Geophysical methods—including electrical resistivity, seismic, and radar—have been increasingly used to characterize the subsurface in a noninvasive manner (e.g. Hubbard and Rubin 2005, Binley *et al* 2015). Geophysics can bridge the gap in sparse wellbore locations by providing high-resolution and

spatially extensive information in a minimally invasive manner. It can image subsurface contaminant plumes (e.g. Johnson *et al* 2010, 2012, Dafflon *et al* 2011), as well as map flow and biogeochemical properties that are important for predictive modeling and understanding (e.g. Johnson *et al* 2010, 2012, Dafflon *et al* 2011, Wainwright *et al* 2014). In particular, geophysics has been powerful in identifying biogeochemical hot spots (Wainwright *et al* 2016a). For the watershed or regional scale, airborne geophysics—particularly airborne electromagnetic surveys—was originally developed for mineral exploration, but is now increasingly used for water-resources applications (e.g. Barfod *et al* 2018, Ball *et al* 2020). While a number of different field experiments have demonstrated the feasibility of such subsurface imaging techniques (e.g. Johnson *et al* 2015), real-time monitoring was still challenging until recently. In particular, autonomous electrical resistivity and phase tomography (ERT) monitoring have the potential to achieve rapid and automated detection and identification of changes in the subsurface (Dafflon *et al* 2011, Johnson *et al* 2015).

In parallel, recent advances in remote sensing have revolutionized how we characterize and monitor ecosystem and watershed dynamics. A high-resolution digital elevation model (DEM) derived from the Light Detection and Ranging (LiDAR) method is increasingly available across the USA, which has been applied to understand better the relationship between geomorphology and hydrology (Prancevic and Kirchner 2019), as well as to measure snow depths and snow-water-equivalent (SWE) over a basin scale (Painter *et al* 2016b). Because LiDAR data has been shown to inform near-surface soil properties (Gillin *et al* 2015, Patton *et al* 2018), hydrological connectivity (Jencso *et al* 2009) and biogeochemical hot spots (Duncan *et al* 2013) in the literature, it is being increasingly used in several DOE sites, including the East River (Hubbard *et al* 2018) and Columbia River Corridor Site at Hanford (Chen *et al* 2021). Similarly, hyperspectral remote sensing is used extensively in DOE- and non-DOE supported research to map plant traits (e.g. Asner *et al* 2015), leaf water contents (e.g. Colombo *et al* 2008), leaf chemistry (e.g. Feilhauer *et al* 2015) and other properties, which are also proxies for soil biogeochemistry (e.g. Madritch *et al* 2014). The East River is making use of these methods to characterize a range of properties, including leaf area, wet, and dry weights for leaf sample, strait models developed independently for needle and non-needle leaf species (Chadwick *et al* 2020a, 2020b, 2020c). For capturing the spatiotemporal heterogeneity, time-lapse images from satellites have enabled the monitoring of vegetation dynamics associated with hydrological disturbances (e.g. Wainwright *et al* 2020). The remote sensing images are available at increasingly high resolution and high frequencies (Falco *et al* 2018, Devadoss *et al* 2020).

Representativeness and quality of collected environmental datasets (e.g. groundwater, precipitation) impact accuracy and precision of climate, hydrological, and biogeochemical analyses and predictions. Often, these are incomplete and may contain unspecified or missing entries due to unavoidable reasons, such as unfavorable weather conditions or delays in collecting sensor data. Several statistical and machine learning-based methods have been developed to render Quality Assurance (QA)/Quality Control (QC) and address related issues using datasets at the East River (Dafflon *et al* 2020, Mital *et al* 2020, Faybishenko *et al* 2021, Dwivedi *et al* 2022).

5. Simulation capability developments (HPC, RTM and other advances)

With the evolution of research on the DOE sites, several important advances have been made, and new computational capabilities have been developed that enable a more robust predictive understanding of the complex processes at work. A more detailed description of these processes is presented below.

5.1. Developments in hydrological and reactive transport modeling tools

The need to understand and predict contaminant transport and the export of nutrients and other important constituents has driven the development of much-improved modeling tools for multiphase (variably saturated) hydrological flow, such as TOUGH (Pruess 1987, 2004, Finsterle *et al* 2014, Jung *et al* 2018, Sonnenthal *et al* 2021), STOMP (White and Oostrom 2000, 2003), PFLOTRAN (Hammond *et al* 2007, 2014, Lichtner *et al* 2015), and reactive transport such as Crunch (Steefel *et al* 2005, 2015), ToughReact (Sonnenthal *et al* 2021), PFLOTRAN (Hammond *et al* 2007, 2014, Lichtner *et al* 2015). In turn, the availability of the new and evolving software has made it possible to address many scientific and engineering issues quantitatively at levels that were not possible previously.

5.1.1. Multiphase hydrology software

The multiphase (variably saturated) hydrological flow software developed for the DOE sites has historically been intended primarily for simulation of subsurface processes based on continuum formulations for porous media Darcy flow. Several hydrological codes such as HP1/HPx (Jacques and Šimůnek 2005, Jacques *et al* 2006, 2012, 2013), PHT3D (Appelo and Rolle 2010, Greskowiak *et al* 2010), OpenGeoSys (Kolditz *et al* 2012, Nagel *et al* 2013), MIN3P (Mayer *et al* 2002), and HYDRUS (Šimůnek *et al* 2006, 2008, Šimůnek and van Genuchten 2008) have been developed globally since the late 1980s and early 1990s. However, many of the multiphase codes for variably saturated flow were developed within the

DOE Laboratory system, including the TOUGH family of codes (Finsterle *et al* 2014), PFLOTRAN (Hammond *et al* 2012), STOMP (White *et al* 2012), NUFT (Hao *et al* 2012), FEHM (Zyvoloski *et al* 1997a, 1997b, Dempsey *et al* 2012), and Amanzi–Advanced Terrestrial Simulator (ATS) (Garimella *et al* 2014, Coon *et al* 2016, 2020). Variably saturated flow has been represented by the Richards equation, which assumes a passive gas phase, as well as with a full multiphase treatment in which multiple phases are subject to gradients and allowed to flow. Enhancements over the years have focused on improving the numerical stability of the Richards or multiphase flow simulations, which are notoriously difficult because of the nonlinearity of the problems they address. In addition, implementation of massively parallel computer architectures has been the focus of much of the effort (e.g. Hammond *et al* 2012, Garimella *et al* 2014, Jung *et al* 2017), although the details of this effort are beyond the scope of this review.

In order to extend beyond the subsurface so as to consider integrated surface and subsurface water, as required for analyzing terrestrial systems that consist of watersheds and river basins, much additional effort has been required. DOE-supported researchers have made important advancements to the state of the art in these integrated surface–subsurface hydrological models, with a focus on highly parallel open-source community codes. For example, extensions (Wu *et al* 2021) of PFLOTRAN (Hammond *et al* 2014) to include surface water and recent updates (Kuffour *et al* 2020) to the integrated surface–subsurface hydrology model PARFLOW (Kollet and Maxwell 2008, Kollet *et al* 2010) were supported by DOE projects. The Amanzi–ATS code (Coon *et al* 2016, 2020) was initiated as subsurface flow and reactive transport code (Amanzi) to support work at DOE legacy sites, and then extended as Amanzi–ATS to include surface flow and reactive transport processes to support research across DOE’s network of watershed research testbeds.

The state of the art with respect to reactive transport modeling was summarized recently in Steefel *et al* (2015). In this summary of a special issue on benchmarking of reactive transport software for environmental applications emphasizing continuum formulations, such codes as PHREEQC, TOUGHReact, OpenGeoSys, PFLOTRAN, MIN3P, eStomp, HydroGeochem, and CrunchFlow were described. The first of these benchmarking workshops was supported by DOE-BER, although at a bare-bones level. The codes were shown to be capable of achieving similar or identical results for a range of complex environmental problems that required numerical solutions, i.e. those with nonlinear or coupled processes that are well beyond what can be described with analytical solutions subject to many simplifying assumptions. A significant portion (but not all) of this software development was undertaken

with funding from the DOE, while the applications of the software were motivated in many cases by environmental problems on DOE lands and later by the desire to understand and predict biogeochemical cycling in ecosystems.

The development of biogeochemical reaction capabilities that could be used in reactive transport codes has followed a somewhat separate path. These efforts have followed closely the fundamental scientific work on understanding the role of microbes in surface and subsurface environments, particularly as they contribute to contaminant retardation through their control of the redox environment (NABIR 2006). Thus, the focus in reactive transport development here has been on incorporating arbitrarily complex biogeochemical reaction networks mediated by one or more competing or collaborating microbial communities. The microbial communities mediate rates that depend on the local environmental conditions and thermodynamics and contribute to reaction pathways that affect contaminant sorption (and thus retardation) and mineral dissolution + precipitation, as well as gas exchange. One example of such a biogeochemical reaction network with multiple microbially mediated pathways is shown in figure 4 (Arora *et al* 2016b).

As might be expected for complex environmental problems like those facing the DOE Hanford Site, where several processes come into play over a large spatial extent, massively parallel computing for reactive transport is now playing an important role. Arguably the first parallel code for reactive transport was the MCTracker code described and verified in Yabusaki *et al* (1998). Following this by a few years was the parallel software PFLOTRAN (Hammond *et al* 2012) that built on many of the capabilities found in FLOTRAN. The power of this new code and its application to the real-world environmental problem of U contamination in the Hanford 300 Area was highlighted in Hammond and Lichtner (2010), discussed below (figure 5). As part of the DOE Advanced Simulation Capability for EM (ASCEM) program, the high-performance computing code Amanzi was developed to simulate subsurface flow and transport. This Amanzi framework was later integrated into the ATS code to bring in surface processes like freeze and thaw (Painter *et al* 2016a), overland flow, and river flow (Coon *et al* 2020).

5.1.2. Integrated surface–subsurface reactive transport software

Although spatially distributed models that fully integrate surface and subsurface flow representations are increasingly available (e.g. Paniconi and Putti 2015), extensions to include reactive transport in surface and subsurface water are relatively rare, Amanzi–ATS includes the capability for integrated surface–subsurface transport and accesses mature biogeochemical reaction capabilities in PFLOTRAN or




Figure 4. Example of a biogeochemical reaction network used to simulate processes in the Rifle flood plain (from Arora *et al* 2016b).




Figure 5. Example of a 3D flow and reactive transport simulation of the Hanford 300 Area, with hydrological and geochemical transients driven by changing water stages in the Columbia River (from Hammond and Lichtner 2010). Hammond and Lichtner (2010). John Wiley & Sons. Copyright 2010 by the American Geophysical Union.

CrunchFlow (see Dwivedi *et al* 2016a) through an Application Programming Interface called Alquimia (github.com/LBL-EESA/alquimia-dev). This relatively new reactive transport capability for integrated surface water-groundwater systems is an important complement to observational studies, with significant potential for aiding in the interpretation of field observations.

Reactive transport in the integrated surface water-groundwater system is important for understanding transport and biogeochemical transformation in river corridors. The hyporheic zone often contains biogeochemical hot spots where many biogeochemical transformations occur. However, the open channel is the primary pathway for downstream movement and allows for O₂ exchange with the atmosphere, emphasizing the importance of an integrated modeling capability. Integrated surface-subsurface reactive transport models have great potential for helping understand the field-scale implications of relatively small-scale hydrologic and geochemical processes occurring in the hyporheic zone. However, it is first necessary to address the mismatch between the native spatial and temporal scales of fundamental hydrologic and biogeochemical processes, and the systems that they impact (e.g. watersheds). For example, locally anoxic conditions within an otherwise O₂-rich environment are important for nutrient and trace-metal processing in stream corridors, but bringing that fundamental understanding into field-scale models remains challenging. DOE-funded researchers have recently introduced multiscale models that confront that scale mismatch by using relatively coarse discretization for the channel network, with subgrid models for hyporheic zone transport and reactions. Painter (2018), and Painter (2021) introduced the ADELS (Advection Dispersion equation with Lagrangian Subgrid) model, which associates a one-dimensional advection-reaction subgrid model with each channel grid cell to represent the ensembles of streamlines that are diverted from the channel into the biogeochemically active hyporheic zone before returning to the channel. The computationally challenging simulations are made tractable by formulating the subgrid models in stochastic Lagrangian form, with hyporheic age replacing travel distance. Key model inputs—hyporheic exchange rates and hyporheic lifetime distributions—can be inferred from nonreacting tracer tests (Rathore *et al* 2021) or synthesized data products (Gomez-Velez and Harvey 2014). The ADELS model is implemented in Amanzi-ATS (Jan *et al* 2021). Motivated by similar considerations, Fang *et al* (2020) implemented a multirate transient storage model in PFLOTRAN. In contrast to the advection-based conceptualization of the ADELS model, the multirate transient storage model (mTSM) considers the hyporheic zone to be a collection of well-mixed reactors that are each coupled to the channel but not to each other. The ADELS and

mTSM models have been shown to be mathematically equivalent for nonreacting tracers but yield very different results for reactive transport, because of the different conceptualization of hyporheic exchange (Painter 2021). These model developments clearly demonstrate that field-scale models that honor fine-scale understanding are possible and have significant potential, but the identified differences in field-scale reactive transport also underscore the importance of the observational studies afforded by DOE's network of watershed research testbeds.

5.2. Advances in contaminant transport modeling

Modern reactive transport methods have made significant contributions to the topic of contaminant transport in recent years, and many of these applications are to DOE legacy sites or to sites where DOE contributed funding (e.g. Naturita, Colorado). The contributions can be broadly divided into two important themes:

- (a) The comprehensive and rigorous treatment of geochemistry in contaminant transport models, and
- (b) The role of aquifer heterogeneity in transport and mixing rates.

The second of these topics is not considered further here, since it is primarily the purview of hydrology and transport disciplines and is beyond the scope of this review paper. We focus here on the geochemical and mineralogical aspects of the problem.

The starting point for contaminant hydrogeology has always been the linear distribution coefficient (or K_d) models for sorption, which have the advantage of being easily incorporated into transport equations without rendering them nonlinear. It has also been asserted that they have an advantage in that the data required for the implementation is minimal, or at least less than that required by the multicomponent models discussed below. However, it should be pointed out that the K_d models, while potentially being as simple as the determination of the sorbed and aqueous concentration under a set of environmental conditions, typically require unique constraints for each and every site to which they are applied. In addition, as will be apparent from the discussion below, such K_d models may not even apply to a single site where conditions (temperature, salinity, competing ion concentrations, sorption site density) change over time and space. In other words, there is no real generality in the case of linear distribution coefficients, in contrast to the more rigorous ion exchange and surface complexation models (like that of metals on Fe-hydroxides) that can be applied quite widely.

The use of linear K_d coefficients came under strong attack when it became apparent that ¹³⁷Cs had migrated further than expected as a result of highly saline tank leaks in the Hanford 200 tank farm




Figure 6. Example of ^{137}Cs transport in the Hanford 200 Area tank farm. (A) Effective linear distribution coefficient (K_d) for Cs^+ (from Steefel *et al* 2003). The K_d depends on both the competing cation concentration (Na^+ associated with NaNO_3 in the tank wastes), and on Cs^+ itself because of the presence of at least two sites with different selectivities for Cs^+ versus Na^+ ; (B) Relative migration of nitrate (nonreactive), and Cs^+ at 1 M NaNO_3 and 5 M NaNO_3 . The high Na^+ concentrations in the tank leak explain the enhanced migration and thus weaker-than-expected retardation of ^{137}Cs at the Hanford 200 tanks (from Steefel *et al* 2005).

(Zachara *et al* 2002, Steefel *et al* 2003, 2005). To address this issue, multicomponent ion exchange and surface complexation models were then considered in the context of reactive transport. Below, we briefly review them.

5.2.1. Ion exchange and Cs migration at Hanford 200 Area

The problem of understanding and predicting ^{137}Cs migration was made more complicated as a result of the multiple exchange sites that occur in sediments below the Hanford 200 tanks (Zachara *et al* 2002, Steefel *et al* 2003). In the Hanford case, the effective K_d depends on the Cs^+ concentration in addition to the competing sodium ion (Na^+) concentration (figure 6(A)), since the ion exchange sites with a very high preference for Cs^+ are present in only low concentrations in the sediment. The low concentration of the high affinity exchange sites means that they could be completely filled, resulting in a reliance on relatively lower affinity sites for the ^{137}Cs sorption.

An example 2D reactive transport simulation was presented in Steefel *et al* (2005) to demonstrate the effect of the competing sodium nitrate (NaNO_3) (a major component of the leaking tank fluids) concentration. This simulation was carried out using ion exchange selectivities and site concentrations determined from Hanford bulk sediment (Steefel *et al* 2003). Figure 6(B) shows the transport of the nonreactive nitrate, with the plume extending over the entire domain size considered. The fractional migration of ^{137}Cs at 1 M NaNO_3 is significantly less, in keeping with the expectation that some retardation of this contaminant will occur (although still more than would be the case for ^{137}Cs in a typically dilute soil or vadose zone water). At 5 M NaNO_3 , the migration of the ^{137}Cs is significantly farther, even if still retarded relative to the nitrate. The enhanced ^{137}Cs observed below many of the Hanford 200 tanks can thus be explained largely by a classical ion exchange model that accounts for the competing NaNO_3 concentrations in the plume, and by the elevated concentrations of Cs^+ that exceed the number of high

affinity sites that are available for strong sorption. The model also suggests that as leaking of highly concentrated NaNO_3 tank fluids ceases, further migration of the ^{137}Cs is unlikely. This is because infiltration by more normal rainwater is not capable of displacing the ^{137}Cs presently sorbed on the Hanford sediment. In any case, the use of constant linear K_d to describe the transport fails completely, so either the more rigorous ion exchange models need to be used, or at minimum, a model that accounts for environmental condition-dependent sorption (i.e. a ‘smart K_d ’).

5.2.2. Surface complexation and U migration

Perhaps an even more convincing demonstration of the inadequacy of the classical K_d models is provided by reactive transport analyses of metal and radionuclide migration influenced by surface complexation on Fe-hydroxides. In fact, an entire generation of geochemists has investigated the surface complexation behavior of metals on ferric hydroxides over many years (Dzombak and Morel 1990, Davis *et al* 1998, 2004), but only more recently has the use of surface complexation models been demonstrated convincingly in reactive transport frameworks at the field scale.

Most likely, the first successful demonstration of the use of surface complexation models to describe field-scale reactive transport was presented by Davis and co-workers for the DOE-LM U-contaminated site at Naturita, Colorado (Curtis *et al* 2004, 2006, Davis *et al* 2004). Davis *et al* (2004) used both electrostatic and nonelectrostatic surface complexation models to describe U sorption on the Naturita sediment, but finally settled on the nonelectrostatic models because of their flexibility in treating natural and complex multiminerale sediments (figure 7(A)). In order to match the pH dependence of sorption, in particular, it was necessary to use a three-site SCM consisting of weak, strong, and very strong sites. Figure 7(B) shows the match with the experimental data at a variety of CO_2 partial pressures, an important variable because of the strong competition between U carbonate complexes in solution and the




Figure 7. Example demonstrating inadequacy of K_d (linear sorption) models to describe U transport. (A) K_d values for Naturita sediment as a function of partial pressure of CO_2 , pH, and solid/liquid ratio (from Davis *et al* 2004). (B) Simulated values of field scale dissolved U, alkalinity, and spatially variable K_d at the Naturita site (From Curtis *et al* 2006). (A) Reprinted from Davis *et al* (2004), Copyright (2004), with permission from Elsevier. (B) Curtis *et al* (2006). John Wiley & Sons. Copyright 2006 by the American Geophysical Union.

surface complexes developed on the Naturita sediment surfaces (Hsi and Langmuir 1985, Prikryl *et al* 2001, Davis *et al* 2004). Calcium uranium carbonate complexes in solution can further reduce the sorption of U, especially $\text{Ca}_2\text{UO}_2(\text{CO}_3)_3$ (Bernhard *et al* 2001, Brooks *et al* 2003, Fox *et al* 2006).

Curtis *et al* (2006) used the nonelectrostatic surface complexation model presented in Davis *et al* (2004) to investigate field-scale transport at the Naturita, Colorado Site (figure 7(B)). They demonstrated that the surface complexation model accurately reproduced the available data when combined with the realistic flow and transport parameters for the aquifer. They also demonstrated the inadequacy of a constant K_d model (figure 7(B)).

Many other studies have been carried out that demonstrate the ability of the surface complexation models to describe field-scale behavior, including those at the U-contaminated Rifle Site (Yabusaki *et al* 2007, Zachara *et al* 2013), and at the contaminated Savannah River Site (Arora *et al* 2018).

5.2.3. Modeling of microbially mediated reaction effects on U migration at the Rifle Site


The fundamental studies of biogeochemistry mediated by microbial communities undertaken as part of the DOE research portfolio over the last 10–15 years have contributed to a much improved ability to model redox processes in the subsurface. This is a key development for predicting the dependence of redox-sensitive contaminants like U and Tc in the subsurface. The redox state of the subsurface depends on the classical sequence of electron acceptor-donor microbially mediated pathways from aerobic respiration under oxidizing conditions to methanogenesis under reducing conditions. The redox state of the subsurface can be manipulated with the injection of a suitable electron donor that drives the system towards reducing conditions, as was demonstrated at the Rifle Integrated Field Research Challenge in

western Colorado (Anderson *et al* 2003, Dullies *et al* 2010, Williams *et al* 2011). The engineered biostimulation of indigenous microorganisms like *Geobacter* (see section 4.1) to catalyze the conversion of U^{VI} to a reduced immobile form of U, U^{IV} , was demonstrated convincingly at the field scale, although the longevity of such treatments has remained in question.

In the field experiments undertaken at the Rifle Site, acetate was injected into a series of wells upgradient of a series of monitoring wells (figure 8(A)). However, to simulate the acetate injection and resulting U reduction and immobilization as a result of microbial (primarily *Geobacter*) activity, it proved useful to begin with 1D column experiments (Li *et al* 2009). With this approach, it was possible to show that the reactive transport modeling could capture both the sequence and distribution of redox processes (figure 8(B)).

To capture the combined behavior of the acetate, U, and other electron acceptors/donors in the 2008 field experiment, it was necessary to use a fully 3D model for the site (Yabusaki *et al* 2011). The injection of acetate drives the system to reducing conditions, with the result that the U^{VI} is immobilized as solid U^{IV} , and the ability to capture this behavior (along with associated redox processes) at the field scale is demonstrated convincingly with the simulations as shown in figure 8(C) (Yabusaki *et al* 2011).

Similarly, to understand the biogeochemical behavior of U within the groundwater and aquifer solids at the Oak Ridge Field Research Center, and as part of their Integrated Field Research Challenge, a conceptual model for simulating an emulsified vegetable oil injection was developed and incorporated into the geochemistry code PHREEQC (Tang *et al* 2013a, 2013b). Numerous field experiments and subsequent interpretation of pyrosequencing and qPCR results in conjunction with geochemical data provided the basis for this hybrid microbial-geochemical conceptual model



development (Gihring *et al* 2011). Consistent with the field observations, the model demonstrated the biologically mediated reductive immobilization of U in groundwater with various electron donors and under diverse geochemical conditions (Wu *et al* 2006a, 2006b, Watson *et al* 2013).

6. Hg: a use case to study watershed biogeochemical processes

Hg is a pollutant of global concern (Assessment, UNEP Global Mercury 2019). Hg has a complex environmental cycle that has been significantly altered by anthropogenic activity. Elemental Hg (Hg^0) has a significant vapor pressure and long residence time in the atmosphere, facilitating long-range transport from source areas. Given that Hg mining dates back over 3000 years, there may be no place on the Earth's surface that has not been affected by this altered cycle.

Inorganic Hg has well-documented deleterious health effects, and its transformation in the environment to the more toxic organomercurial compounds (primarily monomethylmercury, MeHg) poses increased risks to human and environmental health. The MeHg measured in environmental samples is the net result of the counteracting processes of Hg methylation and MeHg demethylation. Research into Hg environmental cycling has a long history, yet fundamental questions remain unanswered, particularly with respect to the watershed biogeochemical conditions that promote net MeHg generation. For example, although the gap is closing, research on lotic ecosystems (rivers and


streams) is relatively underrepresented in comparison to lentic (lakes and ponds) ecosystems.

DOE-supported research on watershed-scale biogeochemistry has been conducted using an iterative pattern-to-process approach, in which the interactions and feedbacks among pattern and process were investigated across a range of spatiotemporal scales. These are interpreted to inform our understanding of watershed-scale function. This is being achieved through a combination of long-term observations, targeted event sampling, studies of seasonal and intra-day patterns, and conducting field-informed experiments under controlled conditions in the laboratory. An important study site is EFPC (see section 3.2), which has a well-documented history of Hg contamination that began at the headwaters and affected the creek and floodplain along its entire length (Brooks and Southworth 2011, Loar *et al* 2011).

6.1. Baseflow sampling identifies additional sources of Hg to the creek

Long-term (~10 year) observations made at baseflow along the length of EFPC demonstrate consistent opposing patterns of Hg and MeHg concentration. Total and dissolved Hg (Hg_T and Hg_D , respectively) concentrations decrease from the headwaters downstream. Conversely, total and dissolved MeHg ($MeHg_T$ and $MeHg_D$) concentrations increase with distance downstream. Additionally, the MeHg concentrations show a seasonal dependence, with higher concentrations in spring and summer and lower concentrations in autumn and winter (figure 9).

Discharge in EFPC increases with distance downstream, suggesting that one explanation for the



decreasing Hg_T and Hg_D concentration is that the additional water is diluting it in the channel. Nevertheless, dilution or particle settling cannot account for the concentration changes. In other words, the fact that the flux of Hg increases with distance downstream suggests that legacy sources of Hg in the watershed contribute additional Hg mass to EFPC (Peterson *et al* 2018). These mass-balance-based estimates are supported by natural abundance Hg stable isotope signature studies (Donovan *et al* 2014, Demers *et al* 2018). By comparing Hg stable isotope signatures from different possible source areas and along the creek, hyporheic water was identified as the likely source of the added Hg, and the reduction of Hg^{II} to Hg^0 by both microbially mediated and photochemical mechanisms was indicated.

Two controlling variables on Hg speciation and transport are NOM and dissolved organic matter (DOM), respectively, and total suspended solids (TSS). Much of the Hg transported downstream is associated with suspended particles comprising a mixture of mineral particles, particulate organic detritus, and algae, predominantly diatoms. The nature of the association between Hg and particle surfaces has broader implications with respect to the reactivity and biogeochemical cycling of Hg in the creek ecosystem, including its release from particles and availability for methylation. Synchrotron-based x-ray fluorescence mapping of the suspended particles indicated that the mineral-bound Hg was closely associated with Fe and NOM (Gu *et al* 2014). In conjunction with complementary Fourier-transform infrared spectroscopy (FTIR) analysis, these results suggested NOM-facilitated Hg bonding to Fe oxides, in which Hg was tightly bound to reduced S groups in the NOM, and the NOM was bound to Fe-oxides via its carboxyl and hydroxide functional groups. The diatom-associated Hg was found on the outer cell surface, suggesting

passive sorption onto the cell surface rather than active uptake.

The filter-passing, or dissolved, Hg is strongly complexed by reduced S groups in DOM (Skylberg *et al* 2006). Using ultrahigh resolution Fourier transform ion cyclotron resonance mass spectrometry (FTICR-MS) coupled with electrospray ionization, investigators found that molecules containing multiple N and S atoms dominated the complexation of Hg (Chen *et al* 2017). From both EFPC and Suwanee River, a 2:1 complex ($[C_5HNS_2-Hg-S_2NHC_5]^+$) that dominated Hg complexation was identified in DOM isolates based on both molecular mass and Hg stable isotope distribution. Possible molecular structures of this complex were estimated using density functional theory calculations.

MeHg flux also increases with downstream distance, which is expected given that both concentration and discharge increase. As is the case for EFPC, MeHg is typically not the initial form of Hg entering the ecosystem; rather, it is formed in the environment from an inorganic Hg precursor via a microbially mediated reaction carried out by a diverse group of anaerobic bacteria possessing the *hgcAB* two-gene cluster (Gilmour *et al* 2013, Parks *et al* 2013, Podar *et al* 2015).

Numerous studies have demonstrated that watershed wetland abundance is positively correlated with MeHg concentrations in the water bodies that drain those watersheds (Wentz *et al* 2014). Nevertheless, the EFPC watershed has 3% wetland abundance, but EFPC has MeHg concentrations typical of watersheds with 30% wetland abundance. Like inorganic Hg, MeHg in the water column is either particle associated or complexed with NOM. An overarching goal of the research is to understand where and under what biogeochemical conditions Hg is methylated and MeHg demethylated.




Figure 10. Flood hydrograph from EFPC illustrating (A) creek discharge and specific conductance as a proxy measure of dissolved ions, and (B) dissolved MeHg and DO dynamics before, during, and after the storm event. The specific conductance mirrors the flood hydrograph diluting during higher flow and returning to pre-flood levels approximately one and a half days after the start of the flood. In contrast, DO and MeHg_D do not return to pre-flood conditions until ten days after the flood start (from Riscassi *et al* 2016).

6.2. Flood sampling reveals terrestrial–aquatic connections and MeHg sources

Targeted sampling of multiple storm-driven flood events revealed specific linkages between the terrestrial and aquatic environments. For instance, precipitation events delivered DOM and associated Hg to the creek, but not MeHg or particulate phases. Additionally, the recovery of dissolved solute concentrations to pre-flood levels closely followed the flood hydrograph, except for MeHg_D which lagged significantly behind other solutes. Recovery of MeHg_D concentrations closely followed the re-establishment of the diel cycle of dissolved O_2 concentration (figure 10). Taken together, the results of targeted flood sampling implied that in-stream biological mechanisms are primarily responsible for MeHg generation in EFPC (Riscassi *et al* 2016).


6.3. Seasonal and intraday patterns highlight the importance of algal biofilms

Natural cycles and anthropogenic activity affect stream ecology, water quality, and water quantity on timescales as short as months to hours. For example, the progression of the seasons leads to cyclic variation in creek discharge, water temperature, sunlight intensity, and duration. Well-established diel patterns of dissolved oxygen (DO) concentration and pH are driven by photosynthetic activity (Allan and Castillo

2007). Wastewater treatment plants and hydroelectric dam operations are anthropogenic causes of short-time-scale water quality variations and river discharge.

Do Hg and MeHg concentrations have analogous cyclic variations to things like the daily photocycle, pH, and DO, and if so, how do they compare with other water-quality variations? Multiple sampling campaigns have been conducted to monitor changes in water chemistry over 30 to 48 hours to address this question. Previous research has documented the photochemical demethylation of MeHg and the role that various photosensitizers, including NOM, play in enhancing or inhibiting the reaction (Qian *et al* 2014). Given these earlier results, lower MeHg concentrations were anticipated during the day, driven by photodemethylation. Instead, MeHg concentrations were positively correlated with the daily photocycle increasing during the day with overnight minima. While photodemethylation may have been occurring, the rate of MeHg production outpaced that of all demethylation reactions, resulting in net positive MeHg production (Brooks *et al* 2021).

Long-term observations demonstrated seasonally dependent watershed sources of MeHg to EFPC. Targeted flood event sampling pointed to in-stream as opposed to out-of-stream sources of MeHg, and



that the recovery of MeHg concentrations parallels the recovery of diel patterns in DO. Diel MeHg concentration variations are closely correlated with the daily photocycle. Together, these observations led us to hypothesize that periphyton biofilms are a source of MeHg in EFPC.

6.4. Field-informed laboratory experiments identify a dominant source of MeHg

Periphyton, or benthic algae, biofilms are complex assemblages comprising biotic (e.g. algae, fungi, bacteria, and archaea, insect larvae), abiotic (mineral grains, entrained sediment particles), and detrital (coarse and fine particulate organic matter) components often embedded in an extracellular gel-like matrix. These biofilms are in-stream, and their photosynthetic inhabitants produce O₂ in a pattern driven by the daily photocycle. During floods, the biofilms are buried and scoured, disrupting the daily DO cycle, which is gradually re-established as the biofilms return. However, bacterial MeHg production is only carried out by anaerobic bacteria. Specifically, the only known methylators are iron-reducing bacteria, SO₄²⁻-reducing bacteria, fermentative microorganisms, and methanogenic archaea. For Hg methylation to occur in periphyton biofilms, these oxygenic systems must also harbor O₂-free zones where these anaerobic microorganisms can be active.

Chemical gradients within intact periphyton biofilms collected from EFPC were studied in the lab using microelectrode voltammetry. Using this technique, concentrations of DO, Fe^{II}, and S^{-II} were measured vertically from the periphyton-water interface down into the biofilm. Over a vertical distance of a few millimeters, conditions progressed from O₂-saturated to iron-reducing and SO₄²⁻-reducing. The steep biogeochemical gradients within the biofilms were consistent with the biofilms harboring conditions conducive to MeHg production.

To further explore the role of periphyton biofilms in Hg cycling, biofilms were grown in the creek, returned to the lab, and amended with enriched stable isotopes of inorganic ²⁰¹Hg and Me²⁰²Hg (Olsen *et al* 2016, Schwartz *et al* 2019). The production of isotopically labeled Me²⁰¹Hg and loss of Me²⁰²Hg was monitored over time as measures of Hg methylation and MeHg demethylation, respectively (figure 11). Experiments were conducted in each of the four seasons. Although the biofilms were a net positive source of MeHg during most incubations, MeHg production was greater during spring and summer and for samples collected downstream versus upstream. Both results are consistent with the long-term field observations that show higher concentrations in those seasons and increased MeHg flux with downstream distance. Biofilms grown or incubated in the dark had substantially lower MeHg production, and in some cases, were net demethylating, which is consistent with the diel sampling in which MeHg concentrations increase during the day and decrease at night. Finally, biofilm structure was important to the measured activity and function of these communities; in biofilms with disrupted structure, methylation potentials decreased by 50%.

High-throughput amplicon sequencing has been applied to methylating periphyton communities to understand better the complex interactions that occur within them (Carrell *et al* 2021). Bacterial and archaeal richness was lower in summer relative to biofilms in the autumn, while fungal richness showed the opposite pattern. Interestingly, Hg methylation potential correlated with numerous bacterial families that do not contain the Hg-methylation genes *hgcAB*, suggesting that overall microbiome structure and relationships within periphyton communities influence rates of Hg transformation.

Hg and MeHg in contact with periphyton are subject to other reactions in addition to methylation and demethylation, including sorption, desorption,

and, in the case of Hg^{II} , reduction to Hg^0 and its reoxidation. These ancillary reactions were measured, and a novel model was created—the transient availability model (TAM)—to more accurately describe the periphyton methylation-demethylation results (Olsen *et al* 2018). Consistent with lab incubations data of field-derived samples from EFPC, the TAM reproduced concentration versus time behavior that had previously been thought to be anomalous. Importantly, the TAM predicts faster Hg methylation rates than previous models, with broad implications for net MeHg generation in lotic ecosystems.

Using Hg biogeochemistry as a use case, we demonstrated that long-term observations in the EFPC, a DOE site (see section 3.2, Oak Ridge Reservation Site), coupled with natural abundance stable isotope fractionation, identified sources and locations of Hg entering the creek. Additionally, it identified seasons of the year that supported higher MeHg concentration. Targeted event sampling demonstrated that discrete and short-duration events connected the terrestrial and aquatic ecosystems in the watershed. In other words, these floods constitute hot moments with major implications for material and energy exchange. Event sampling also eliminated the floodplain as a source of MeHg to the creek and focused attention on in-stream sources. High temporal resolution sampling over daily periods produced the unexpected result of increasing MeHg concentration during the day. Controlled laboratory experiments confirmed that periphyton biofilms are a net positive source of MeHg and constitute hot spots of Hg, Fe, and S cycling. Further, biofilms with a higher degree of connectivity among its populations also have higher Hg methylation potential. This combination of approaches buttressed some ideas, but more importantly, highlighted previous misconceptions, identified HSHMs of governing processes, and opened new avenues of discovery to advance further our understanding of these systems that support diverse and vital ecosystem services for the global economy.

7. Evolution of scientific approaches and strategies to tackle scale and complexity

Significant progress has been made in process understanding at the hillslope and field scales, which is relevant in particular for watersheds such as the mountainous East River Site. Yet watershed-scale modeling has remained challenging because of the scale and wide underlying range of heterogeneity in watershed processes and parameters. Hyper-resolution watershed-scale models capable of exploiting high-performance computing are needed to resolve process complexity (level of details needed) and the spatial gradients that typically drive flow. However, these computationally expensive models pose a serious impediment to scientific

progress, requiring abundant CPU hours. The variable resolution mesh or more sophisticated adaptive mesh refinement methods make it possible to focus on functionally dynamic regions compared to more conventional approaches. Efforts are underway to advance watershed-scale modeling using leadership-class computers, big data, and machine learning combined with learning-assisted physics-based simulation tools (e.g. <https://exasheds.org/>). However, the details of these efforts are well beyond the scope of this review. As an alternative, we have developed strategies to tackle scale and complexity over the years. Some examples of these strategies include multiscale, multiphysics, hybrid modeling, incorporating HSHMs, and functional zonation as part of scale-adaptive modeling.


7.1. Multiscale, multiphysics and hybrid modeling

Historically there has been a mismatch between the spatial and temporal scales at which fundamental hydrologic and biogeochemical processes are most readily studied and understood (molecular to laboratory) and those of the ecosystems that they impact (field to watershed/basin). Because of nonlinear process interactions and complex heterogeneity in natural ecosystems, small-scale (e.g. laboratory) studies do not directly translate to field settings. Key reactive transport parameters, including permeability, dispersivity, and reaction-rate coefficients are all well known to be scale dependent and, in some cases, boundary-condition dependent. These challenges have motivated a variety of modeling approaches that are aimed at integrating information across scales and representing ecosystem processes at multiple levels of fidelity and complexity through DOE- and non-DOE supported research.

Multiple approaches have been developed to address this challenge, each with advantages and limitations. A summary of selected approaches used in DOE-supported research is provided here; for broader reviews, see Keyes *et al* (2013) and Scheibe *et al* (2015a).

Multiresolution methods use variable discretizations to capture effects of small-scale processes and properties on large-scale phenomena (e.g. Özgen-Xian *et al* 2020). These methods include adaptive grid refinement (refining grids dynamically to put higher resolution where and when needed); multigrid methods (hierarchical sets of grids used to speed numerical convergence of a fine-grid solution), and nested or spatially varying grids that enhance resolution in specific subdomains (e.g. figure 12).

Upscaling methods use assumptions about the nature of subgrid variability to derive effective parameters valid at the model resolution. These methods include numerical upscaling (direct solution of subgrid domains to derive approximate grid-scale parameters), various forms of homogenization and



volume averaging, and perturbation solutions to stochastic partial differential equations.

Hybrid multiscale methods are similar to multiresolution methods. However, they use fundamentally different models in selected subdomains (for example, pore-scale simulations in portions of a domain where mixing-limited precipitation occurs; Scheibe *et al* 2015b) and are sometimes referred to as ‘adaptive physics’ models. These include hierarchical methods (e.g. the heterogeneous multiscale method; E *et al* 2003) in which overlapping domains exchange information and concurrent methods (e.g. mortar methods; Balhoff *et al* 2008), within which distinct subdomains pass information at domain boundaries.

7.2. HSHM: a construct to cope with intense biogeochemical activity

The HSHM construct provides the basis for developing a robust predictive understanding of how contaminant behavior in subsurface heterogeneous media varies across spatiotemporal scales. Subsurface heterogeneities are sites of intense biogeochemical activity and are critical to watershed function. Recent studies have highlighted in greater detail the roles of the naturally reduced zones (or ‘redox heterogeneities’) on biogeochemical function. For example, heterogeneity caused by clay lenses in high permeability soils provides opportunities to sorb contaminants in place (Campbell *et al* 2012, Janot *et al* 2016), while fracture networks can lead to rapid infiltration of contaminants to groundwater (e.g. Berkowitz 2002, Arora *et al* 2019a). Thus, high contaminant concentrations can be spatially diffused across large scales or concentrated at higher rates in specific areas,

thereby forming contaminant hot spots. Similarly, these concentrations can be temporally diffused or more concentrated during specific times, thereby exhibiting contaminant hot moments. In addition, hydrological perturbations in transiently saturated zones drive changes in soil microbial communities, triggering a range of geochemical responses, thereby forming dynamic HSHMs. Figure 12 illustrates static hot spots (i.e. fixed in space) and dynamic hot spots (i.e. migrating in space).

Although the HSHM construct simplifies the characterization of complex heterogeneous systems and time scales at which biogeochemical processes occur—by subdividing the region and time into a finite number of relatively homogeneous units—it lacks a quantitative definition. Moreover, the HSHM construct is underutilized because of its dynamic nature. In response to that, Bernhardt *et al* (2017) argued that the temporal dynamics of a putative hot spot are a fundamental trait that should be used in their description and, therefore, stressed the merger of HSHMs into a single descriptive term ‘ecosystem control points.’ This definition indicates that knowledge of the rate distributions has more relevance than knowledge of maximum rates. Partly supported by DOE, Bernhardt *et al* (2017) suggested four distinct categories of ecosystem control points: (a) permanent control points, experiencing high rates of biogeochemical activity relative to their proximity (e.g. floodplain), (b) activated control points, where higher biogeochemical activity is set off when certain substrates are delivered (e.g. low-lying topographic positions), (c) export control points that accumulate reactants until some hydraulic or diffusion threshold

Table 2. Examples of HSHMs in the context of ecosystem control points.

Examples	Study	Type of ecosystem control point
Storm events constituted hot moments of Hg and MeHg in EFPC and U discharge in Tims Branch, resulting in orders-of-magnitude higher concentrations than observed during baseflow conditions	Batson <i>et al</i> (1996), Riscassi <i>et al</i> (2016)	Activated control points
Seasonal river stage fluctuations of more than 2 m resulted in U hot moments at the Hanford Site by promoting water intrusion and increasing residence times	Zachara <i>et al</i> (2020)	Activated control points
Naturally reduced zones or zones enriched with Fe and S minerals in sediments adjacent to the riverbanks at the Rifle Site constituted hot spots of C, N, S, and U	Campbell <i>et al</i> (2012), Janot <i>et al</i> (2016), Arora <i>et al</i> (2016a), Wainwright <i>et al</i> (2016b), Boye <i>et al</i> (2017), Noël <i>et al</i> (2017b), Dwivedi <i>et al</i> (2018a)	Permanent control points
Seasonal meltwater-driven water-table rise at the Rifle Site creates short-lived carbonate solutes (calcium ions) for 2-week, but intense oxidizing hot moments, during which the majority of annual U oxidation occurs	Lezama-Pacheco <i>et al</i> (2015)	Activated control points
Microtopographic features such as low-centered polygons act as CH ₄ hot spots in arctic tundra environments and have a significant impact on ecosystem-scale C fluxes	Vaughn <i>et al</i> (2016), Arora <i>et al</i> (2019c)	Activated control points
Meander acts as a sink for organic and inorganic C as well as Fe during the extended baseflow and high-water conditions	Dwivedi <i>et al</i> (2018b)	Export control points
Microtopographic features such as gullies at the East River Site exert significant impacts on redox processes in the hyporheic zone and thereby controlling subsurface geochemical exports of Fe and C	Dwivedi <i>et al</i> (2018b), Rogers <i>et al</i> (2021)	Transport control points

is not surpassed (e.g. geochemical exports through meanders), and (d) transport control points having extremely high transport capability without their direct intervention in exhibiting high biogeochemical activity (e.g. preferential flow paths). Table 2 presents some examples of HSHMs from DOE research in the backdrop of ecosystem control points.

7.2.1. Impact of HSHMs on biogeochemistry of legacy contaminants and nutrients

DOE investigations at former U ore processing sites within the Upper Colorado River Basin show the importance of HSHMs on legacy contamination (e.g. U) and nutrient cycling (e.g. Fe, S). Clayey lenses containing elevated organic C are common within coarse alluvium. These lenses host intense reducing activity (i.e. hot spots) and sustain halos of reducing conditions within the surrounding coarser aquifer sediments (Kumar *et al* 2020a). Precipitation of iron monosulfides within these lenses and reductive attenuation of U^{VI} to U^{IV} cause S and U to become enriched to levels 25 and 200 fold higher, respectively, than surrounding sediments (Janot *et al* 2016, Noël *et al* 2017a, 2017b, Lefebvre *et al* 2019). As and non-redox active species such as Zn have also been

shown to accumulate under these conditions through multiple chemical mechanisms involving association with reduced S species, as well as precipitation of double hydroxides (Kumar *et al* 2020a, Engel *et al* 2021). When clayey sulfidic lenses reside within the zone of annual water-table excursion, exposure to O₂ perfusion during summer drought conditions can cause SOM-complexed U^{IV} species to be oxidized to U^{VI}, providing a mechanism of release and contamination of groundwater. Desiccation of sediments during summer drought leads to precipitation of uranyl carbonates and silicates (Noël *et al* 2019).

Although the relationship between hydrological change and microbial function is centrally important for predicting watershed function, only a few studies have explored the microbial ecology of dynamic HSHMs. Cardarelli *et al* (2020) examined the microbial ecology of N-cycling archaea within transiently saturated and reduced zones at five contaminated DOE legacy floodplain sites (Shiprock in New Mexico, Rifle, Grand Junction, and Naturita in Colorado, and Riverton in Wyoming), which spans a 900 km N–S transect of the Upper Colorado River Basin. This study showed that ammonia-oxidizing ‘ecotypes’ are strongly influenced by water table

position, the location of reducing zones, or both. This study revealed that archaeal ammonia oxidation predominates within the terrestrial subsurface beyond a meter belowground. Recently, (Tolar *et al* 2020) characterized microbial community diversity at the U-contaminated Riverton, Wyoming, legacy across transiently reduced zones. These results showed that microbial communities are surprisingly stable, despite robust flood-to-drought fluctuation and accompanying redox inversion. These findings were interpreted to indicate that microorganisms oscillate between 'active' and 'dormant' states, contingent upon prevailing environmental conditions, and their metabolic products play a more significant role in contaminant mobility than the organisms themselves. Taken together, these results indicate that microbial community responses to drought and flood are complex and require more study.

7.3. Functional zonation

Modeling eco-hydro-biogeochemical processes requires the characterization of the bedrock-to-canopy properties across watersheds. The tremendous heterogeneity of these properties is often a significant obstacle toward scaling modeling efforts from the hillslope and floodplain scales to a larger scale. Although various spatially extensive characterization techniques such as remote sensing and geophysics are available, it is still challenging to estimate each property distribution individually over the watershed scale, particularly soil and subsurface properties in the subsurface.

A functional zonation approach holds tremendous potential for tractably estimating individual property distributions over the watershed scale. In this approach, zones or spatial units within a landscape with unique distributions of multiple properties relative to neighboring regions can be identified using machine-learning-based spatial clustering approaches (e.g. figure 12). The zonation approach essentially builds on a hydrofacies approach, which has been used in groundwater hydrology to define hydrological parameters in each facies (e.g. Fogg *et al* 1998, Klingbeil *et al* 1999). The reactive facies (Sassen *et al* 2012, Wainwright *et al* 2014) defined both hydrological and geochemical properties in each facies and stochastically mapped the reactive facies based on multiple geophysical data. Bea *et al* (2013) documented the value of incorporating estimated reactive facies information within a reactive transport model to improve predictions of long-term U plume transport. These concepts have proved valuable in identifying zones and hot spots of biogeochemical activity (Flores Orozco *et al* 2014, 2018, Wainwright *et al* 2015).

More recently, the availability of enhanced remote-sensing (including hyperspectral) products

and advanced machine-learning approaches has led to a paradigm shift in the zonation approach and demonstrated possibilities for characterizing the bedrock-to-canopy properties across watersheds. With this new paradigm shift, the zonation approach harnesses a suite of remote sensing datasets for identifying and distributing zones relevant to biogeochemistry and/or ecosystem functioning. In addition, the zonation approach takes advantage of the current scientific understanding of critical zone processes and insights into water and nutrient cycling and their exports within a watershed and its different compartments, particularly through the Critical Zone Observatory (CZO) network (Brantley *et al* 2017).

Next, we briefly describe some prominent applications of the zonation approach at DOE sites. Hubbard *et al* (2013) and Wainwright *et al* (2015) used geophysical and remote sensing datasets to identify zones representing the heterogeneity of key properties and ecosystem functions—such as soil properties and C fluxes—in Arctic tundra. Devadoss *et al* (2020) defined the ecosystem functioning zones (each of which has distinct time series signatures of soil moisture and plant dynamics) and investigated the relationship among soil, plant, and snow dynamics. Wainwright *et al* (2016b) subsequently incorporated U reactive facies (e.g. Bea *et al* 2013) into the zonation approach to identify biogeochemical hot spots (e.g. naturally reduced sediments) based on electrical geophysical methods at the Rifle Site. Several additional modeling studies at the Rifle Site represented naturally reduced sediments in their models to investigate the HSHMs of N, C, Fe, and DO and quantified their subsurface geochemical exports (e.g. Arora *et al* 2016b, Yabusaki *et al* 2017, Dwivedi *et al* 2018a).

Recent investigations have been focused on examining how properties associated with specific functional zones exert controls on water and N exports in response to snow dynamics and contribute to the aggregated watershed concentration–discharge signature in mountainous regions like the East River (Hubbard *et al* 2018). A suite of remotely sensed bedrock-through-canopy watershed data layers and machine-learning approaches were utilized at the East River to identify watershed zonation. The zonation approach explored the sensitivity of different identified zones to foresummer drought (Wainwright *et al* 2022). Zonation is also powerful for understanding the co-variability of above- and belowground processes, as well as identifying relevant field experimental sites throughout the domain. For example, research at the Hanford Site is using river channel morphology zonation (hydromorphic units, Ren *et al* (2021)) as a framework for a transferable understanding of river–groundwater exchange fluxes and transit times.

8. Current ecosystem science and legacy contamination

Current ecosystem science comprehensively aims to couple and model ecological, hydrological, biogeochemical processes across the critical zone, ranging from the bedrock to the canopy. Water security challenges have motivated current ecosystem science, which has continued to emerge and expand well beyond traditional contaminant remediation-type studies. Recent severe weather events and hydro-climatic conditions have only added to these challenges, raising important concerns about geochemical exports of metals, C, N, and nutrients to the environment. In past assessments, the contribution to the riverine budgets of C and N were largely neglected, as part of the monitored natural attenuation strategy exclusively focused on stabilizing contaminants in place. More recently, an increased focus has been placed on constraining the exchanges and fates of different forms of C and N in river-floodplain settings, because of their important role in driving biogeochemical interactions with contaminants, and the potential of increased fluxes under changing precipitation regimes. Several studies have emphasized strong interlinkages between subsurface legacy contamination and hydrologically driven biogeochemical exports of metals, C, N, and nutrients in watersheds (e.g. Stucker *et al* 2013, Zhang *et al* 2018, Kumar *et al* 2020b).

To illustrate this further, riparian zones and wetlands, where the contaminated groundwater interfaces with surface water, play a significant role in transforming contaminants because the presence of plant roots may facilitate the sequestration of trace metals. Large biogeochemical gradients exist in wetland ecosystems influenced by the primary production activities of plants that release large quantities of labile organic matter, and by the transport of O₂ in roots to maintain aerobic respiration. Plant roots also modulate mineralogy and promote contaminant immobilization through altering contaminant binding environment (e.g. Kaplan *et al* 2016). Several studies have explored *in situ* immobilization of contaminants for remediation at several DOE sites (e.g. Szecsody *et al* 1998, Dai *et al* 2002, Pawloski *et al* 2009, Zachara *et al* 2013). Notably, the long-term sustainability of *in situ* reductive bioimmobilization of Cr^{VI} was examined under different biogeochemical regimes at the Hanford 100 Area (e.g. Faybishenko *et al* 2008, Beller *et al* 2014, Varadharajan *et al* 2017). Alternatively, plants can offer an effective solution for the removal of hexavalent Cr by transforming it to less mobile, reduced Cr species as shown in the literature (Xu and Jaffé 2006). Further, O₂ diffused from the wetland plant roots into their surrounding sediments can result in distinctive Fe plaques that precipitate on the surface of roots (Blute *et al* 2004), which may consequently facilitate active Fe cycling

by stimulating Fe-reducing bacteria such as *Geobacter* spp. in anaerobic wetland sediments of the Savannah River Site (Chang *et al* 2014). The enhanced Fe cycling can impact U attenuation by accumulating U near the plant roots in more oxidized states (U^{VI} or U^V), which are less susceptible to remobilization during episodic reoxidation events compared to U^{IV}, as observed at the Savannah River Site (Gilson *et al* 2015, Kaplan *et al* 2016).

Wetlands and river corridors experience episodic oxidation events associated with wetting and drying cycles Noël *et al* (2017a), Tolar *et al* (2020). Such cycling may facilitate the ongoing formation of fresh Fe oxyhydroxide, which can sorb or co-precipitate with U. The flow and transport of water links vegetation, soil, and contaminant transformations at the contaminated sites. The fraction of soil pore space that is water-filled vs. air-filled, i.e. the soil saturation, determines the rate at which water and O₂ can be transmitted to meet the transpiration and respiration demands of plant roots and aerobic soil microbes, thereby influencing soil redox potential (Yan *et al* 2018). Aerobic microsites may form around the roots of wetland plants growing in saturated soils that are otherwise anoxic, leading to reaction hot spots. Redox reactions and cycling of redox-sensitive elements can alter the solubility and availability of key nutrients and metals, shifting microbial metabolism from largely oxidative reactions to methanogenesis, denitrification, and other less energetically efficient processes (Megonigal and Neubauer 2019, Cardarelli *et al* 2020). In turn, the history of water availability and rooting zone redox potential have direct and indirect effects on vegetation's distribution, structure, physiological state, and health.

9. Global water security issues and possible solutions through DOE-supported science

The existence of contaminants and the presence of high levels of nutrients in the environment is a major threat to water security. The problem of freshwater contamination is more pressing than ever, because most of the world's prominent rivers, supporting several million people, are under threat. For instance, India's Ganges River is considered one of the most polluted rivers globally: it contains many chemicals, including toxic Cr (Tripathi *et al* 2016). A major drinking water source in China, the Yellow River, the second-longest in Asia and the sixth-longest river globally, has been suffering from severe pollution and is now on the verge of becoming unusable even for agricultural or industrial use (Brown and Halweil 1998, Lu *et al* 2018). The Citarum, the longest and largest river in West Java, Indonesia, faces extreme pollution from several contaminants, including Hg and As. The Hg concentrations in some parts of the river are as much as 100 times higher than levels

considered acceptable by the EPA (maximum contaminant level is 2 ppb) (Paddison 2016). As another example, every year the Mississippi River in the USA receives millions of pounds of toxic waste, poisonous chemicals (e.g. As), and excess nutrients (e.g. N), thereby turning the river's mouth into a dead zone (Dodds 2006). The Colorado River and its tributaries, supporting over 36 million people in seven states in the western USA, have also experienced contamination from a variety of chemicals, including U and As (Gross 2017). Groundwater contamination by As in Bangladesh represents one of the largest mass poisonings in history, with over a million people exposed to elevated As levels in their drinking water (Islam *et al* 2010).

An accurate accounting of the fate and transport of these chemicals in the environment is key to ensuring water security. Many insights developed from several decades of research work at DOE sites described in this paper can be used globally to examine the fate and transport of various contaminants and nutrient cycling. Capabilities developed as part of DOE research include hydrological and reactive transport modeling tools. These tools can comprehensively address water security problems and are transferable across DOE sites and worldwide. For example, several DOE codes such as PFLOTRAN and CRUNCH are currently being applied for various environmental applications globally (e.g. Alt-Epping *et al* 2015, Knabe *et al* 2021).

Finally, current ecosystem science is evolving through emerging and promising developments that include the use of artificial intelligence methods, such as machine learning to support zone identification, the association of system properties and functions with zonal types, HSHM detection, and the development of macroscale reduced-complexity models (Painter 2018, 2021, Fang *et al* 2020). Collaborations among hydrologists, geochemists, and microbiologists have led to advances in the use of increasingly abundant data from molecular biology studies ('omics') to inform reactive transport models that are beginning to mature (Meile and Scheibe 2019). Multiscale heterogeneity and nonlinear process interactions in natural systems continue to challenge our predictive understanding, but these enhanced models and other recent advances offer promise for the future.

10. Summary, conclusions, and future directions

Within the DOE's Office of Science, the BER and other related DOE programs (e.g. ASCR, EM, ER) have contributed significantly to environmental sciences' progress since the late 1980s. They have addressed several challenging subsurface problems, including treating radioactive waste (e.g. U, Pu) and heavy metals and metalloids (e.g. Cr, Hg, As).

These efforts have provided transferable insights into process understanding, designed novel monitoring strategies to collect valuable field information, developed robust predictive capabilities, and devised scale-aware approaches to tackle underlying processes and the complexity resulting from the ecosystems' scales and magnitudes. To document these scientific advances that are generalizable and applicable to a range of water security problems worldwide, we have synthesized research activities conducted at representative DOE sites and testbeds, including the Savannah River Site in South Carolina, Oak Ridge Reservation in Tennessee, Hanford in Washington, Nevada National Security Site in Nevada, Riverton in Wyoming, and Rifle and East River in Colorado, over the last two decades. In particular, we have described the progress made in subsurface environmental sciences through DOE programs. To demonstrate the range of topics that have been studied in the environmental sciences through DOE-supported research, we show a word cloud highlighting key contributions (figure 13), including keywords such as 'subsurface, environment, contaminant, redox, reactive transport modeling, and watershed'.

Efforts through DOE-supported research have marked several breakthroughs, such as isolation of archaea from groundwater and expansion of the microbial 'tree of life' using genomic methods in the subsurface microbiology; understanding the geochemical and biogeochemical behavior of S, Fe, Mn, U, Cr, Hg, C, and N in redox dynamic environments; and generating a robust predictive understanding of the relevant hydro-biogeochemistry and eco-hydro-biogeochemistry through novel field measurements and simulations. These findings have led the way in predicting the behavior and formation of HSHMs in the environment. Multiscale, multiphysics, and hybrid models have provided a framework for rigorous reactive flow and transport simulation capabilities. These modeling capabilities have the potential to telescope into watershed subsystems to simulate fine-scale detail, taking advantage of advanced meshing workflow, machine learning, and functional zonation concepts.

We also discussed a paradigm shift in the ways that environmental science was done in the early 2010s. Until then, investigations of small-scale (molecular to the millimeter) processes had been a primary focus. This approach was inadequate for addressing scaling behavior in the presence of a range of complex, coupled, nonlinear processes and a wide range of landscape heterogeneity. A holistic system science approach evolved to address these concerns, standing on the shoulders of active contaminant remediation-type studies. Since then, the DOE has supported the system science approach through sustained investments and bringing in multidisciplinary, multi-institutional teams together to unravel the intricacies and interconnectedness of system components that

diverse multiscale data into models for near-real-time prediction, the rapid identification of system-tipping-point precursors, and the development of models that inform the real-time optimization of autonomous sensing systems—from watershed to water basin to continental scales (Hubbard *et al* 2020). Advanced machine learning approaches have the potential to provide actionable intelligence, which will enable efficient data management, smooth workflows, and interdisciplinary collaborations, all contributing to the scientific advancements needed to tackle water security problems in the future.

Data availability statement

No new data were created or analysed in this study.

Acknowledgments

The work highlighted here was chosen because it was overwhelmingly supported by the DOE and would not have existed otherwise. However, the authors did not specifically require that each cited manuscript have a specific DOE funding source acknowledgment. The efforts of DD, CIS, BA, JB, SSH, PN, HMW, and KHW were supported by the Watershed Function Scientific Focus Area (SFA) funded by the US Department of Energy, Office of Science, Office of Biological and Environmental Research under Award Number DE-AC02-05CH11231. DD and CIS also acknowledge support from the ExaSheds Project at Lawrence Berkeley National Laboratory funded by the United States Department of Energy, Office of Science, Biological and Environmental Research under Contract No. DE-AC02-05CH11231. HMW also acknowledges support from the Department of Energy, Office of Environmental Management, ALTEMIS—Advanced Long-Term Environmental Monitoring Systems project. This research used resources of the National Energy Research Scientific Computing Center (NERSC), a United States Department of Energy Office of Science User Facility located at Lawrence Berkeley National Laboratory, operated under Contract No. DE-AC02-05CH11231. The efforts of MIB, DIK, KMK, and EJO were supported by the Wetlands Hydrobiogeochemistry Scientific Focus Area (SFA) at Argonne National Laboratory, which is supported by the Earth and Environmental System Science Program, Office of Biological and Environmental Research (BER), Office of Science, US Department of Energy (DOE), under Contract DE-AC02-06CH11357. JB's effort was supported by the SLAC Floodplain Hydro-Biogeochemistry SFA, funded by the US Department of Energy, Office of Biological and Environmental Research, Earth and Environmental Systems Sciences Division. SLAC National Accelerator Laboratory is supported by the US Department of Energy, Office of Science, Office of Basic Energy Sciences under Contract

No. DE-AC02-76SF00515. The efforts of EP, SP, and SB were sponsored in part by the Office of Biological and Environmental Research within the Office of Science of the US Department of Energy (DOE), as part of the Hg Science Focus Area (Critical Interfaces SFA) and IDEAS-Watersheds projects at the Oak Ridge National Laboratory (ORNL). The DOE will provide public access to these results of federally sponsored research in accordance with the DOE Public Access Plan (<http://energy.gov/downloads/doe-public-access-plan>). ORNL is managed by UT-Battelle, LLC under Contract No. DE-AC05-00OR22725 with DOE. XC and TS were supported by the United States Department of Energy, Office of Science, Office of Biological and Environmental Research, Environmental System Science (ESS) Program through the River Corridor Scientific Focus Area project at Pacific Northwest National Laboratory. MZ's contribution was performed with funding from the Department of Energy, Office of Science, Biological and Environmental Research, Subsurface Biogeochemical Research program (SCW1053) and performed under the auspices of the US Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344. We thank Diana Swantek (LBNL) and Adam Malin (ORNL) for assistance with preparing figures 1, 3, and 8(A). We express our gratitude to Dan Hawks (LBNL) for helping with technical editing. Finally, we thank the anonymous reviewer(s) for their insightful comment that helped improve the manuscript.

ORCID iDs

Dipankar Dwivedi  <https://orcid.org/0000-0003-1788-1900>

Carl I Steefel  <https://orcid.org/0000-0002-2802-611X>

Bhavna Arora  <https://orcid.org/0000-0001-7841-886X>

Jill Banfield  <https://orcid.org/0000-0001-8203-8771>

John Bargar  <https://orcid.org/0000-0001-9303-4901>

Maxim I Boyanov  <https://orcid.org/0000-0001-8758-5248>

Scott C Brooks  <https://orcid.org/0000-0002-8437-9788>

Xingyuan Chen  <https://orcid.org/0000-0003-1928-5555>

Susan S Hubbard  <https://orcid.org/0000-0003-2966-5631>

Dan Kaplan  <https://orcid.org/0000-0002-1019-8386>

Kenneth M Kemner  <https://orcid.org/0000-0002-1409-8542>

Peter S Nico  <https://orcid.org/0000-0002-4180-9397>

Edward J O'Loughlin  <https://orcid.org/0000-0003-1607-9529>
 Eric M Pierce  <https://orcid.org/0000-0002-4951-1931>
 Scott L Painter  <https://orcid.org/0000-0002-0901-6987>
 Timothy D Scheibe  <https://orcid.org/0000-0002-8864-5772>
 Haruko M Wainwright  <https://orcid.org/0000-0002-2140-6072>
 Kenneth H Williams  <https://orcid.org/0000-0002-3568-1155>
 Mavrik Zavarin  <https://orcid.org/0000-0002-2426-6706>

References

- Abdelouas A, Lutze W and Nuttall H 1999 Oxidative dissolution of uraninite precipitated on Navajo sandstone *J. Contaminant Hydrol.* **36** 353–75
- Al-Shayeb B et al 2020 Clades of huge phages from across Earth's ecosystems *Nature* **578** 425–31
- Allan J D and Castillo M M 2007 *Stream Ecology: Structure and Function of Running Waters* (Berlin: Springer Science & Business Media)
- Allard S and Gallard H 2013 A biotic formation of methyl iodide on synthetic birnessite: a mechanistic study *Sci. Total Environ.* **463** 169–75
- Alley W M, Healy R W, LaBaugh J W and Reilly T E 2002 Flow and storage in groundwater systems *Science* **296** 1985–90
- Alt-Epping P, Tournassat C, Rasouli P, Steefel C, Mayer K, Jenni A, Mäder U, Sengor S and Fernández R 2015 Benchmark reactive transport simulations of a column experiment in compacted bentonite with multispecies diffusion and explicit treatment of electrostatic effects *Computat. Geosci.* **19** 535–50
- Anantharaman K et al 2016 Thousands of microbial genomes shed light on interconnected biogeochemical processes in an aquifer system *Nat. Commun.* **7** 1–11
- Anderson R T et al 2003 Stimulating the *in situ* activity of *Geobacter* species to remove uranium from the groundwater of a uranium-contaminated aquifer *Appl. Environ. Microbiol.* **69** 5884–91
- Appelo C and Rolle M 2010 PHT3D: a reactive multicomponent transport model for saturated porous media *Ground Water* **48** 627–32
- Arora B et al 2016b Influence of hydrological, biogeochemical and temperature transients on subsurface carbon fluxes in a flood plain environment *Biogeochemistry* **127** 367–96
- Arora B, Burrus M, Newcomer M, Steefel C I, Carroll R W, Dwivedi D, Dong W, Williams K H and Hubbard S S 2020 Differential CQ analysis: a new approach to inferring lateral transport and hydrologic transients within multiple reaches of a mountainous headwater catchment *Front. Water* **2** 24
- Arora B, Davis J A, Spycher N F, Dong W and Wainwright H M 2018 Comparison of electrostatic and non-electrostatic models for U^{VI} sorption on aquifer sediments *Groundwater* **56** 73–86
- Arora B, Dwivedi D, Faybishenko B, Jana R B and Wainwright H M 2019a Understanding and predicting vadose zone processes *Rev. Mineral. Geochem.* **85** 303–28
- Arora B, Dwivedi D, Hubbard S S, Steefel C I and Williams K H 2016a Identifying geochemical hot moments and their controls on a contaminated river floodplain system using wavelet and entropy approaches *Environ. Model. Softw.* **85** 27–41
- Arora B, Ireson A, Bouteiller C L, Hector B, Ali G, Stegen J, Wymore A, Sullivan P L, and Groh J 2019b Toward an international critical zone network-of-networks for the next generation through shared science, tools, data, and philosophy
- Arora B, Wainwright H M, Dwivedi D, Vaughn L J, Curtis J B, Torn M S, Dafflon B and Hubbard S S 2019c Evaluating temporal controls on greenhouse gas (GHG) fluxes in an Arctic tundra environment: an entropy-based approach *Sci. Total Environ.* **649** 284–99
- Asner G P, Martin R E, Anderson C B and Knapp D E 2015 Quantifying forest canopy traits: imaging spectroscopy versus field survey *Remote Sens. Environ.* **158** 15–27
- Assessment, UNEP Global Mercury 2019 *UN Environment Programme* (Geneva: Chemicals and Health Branch)
- Balhoff M T, Thomas S G and Wheeler M F 2008 Mortar coupling and upscaling of pore-scale models *Computat. Geosci.* **12** 15–27
- Ball L B, Davis T, Minsley B J, Gillespie J M and Landon M K 2020 Probabilistic categorical groundwater salinity mapping from airborne electromagnetic data adjacent to California's Lost Hills and Belridge oil fields *Water Resour. Res.* **56** e2019WR026273
- Barfod A A, Möller I, Christiansen A V, Høyer A-S, Hoffmann J, Straubhaar J and Caers J 2018 Hydrostratigraphic modeling using multiple-point statistics and airborne transient electromagnetic methods *Hydrol. Earth Syst. Sci.* **22** 3351–73
- Bargar J R et al 2013 Uranium redox transition pathways in acetate-amended sediments *Proc. Natl Acad. Sci.* **110** 4506–11
- Bargar J R, Fuller C C, Marcus M A, Brearley A J, De la Rosa M P, Webb S M and Caldwell W A 2009 Structural characterization of terrestrial microbial Mn oxides from Pinal Creek, AZ *Geochim. Cosmochim. Acta* **73** 889–910
- Bargar J R, Reitmeier R, Lenhart J J and Davis J A 2000 Characterization of U^{VI}-carbonate ternary complexes on hematite: EXAFS and electrophoretic mobility measurements *Geochim. Cosmochim. Acta* **64** 2737–49
- Batson V L, Bertsch P, and Herbert B 1996 Transport of anthropogenic uranium from sediments to surface waters during episodic storm events *Technical Report* (Wiley Online Library).
- Bea S A, Wainwright H, Spycher N, Faybishenko B, Hubbard S S and Denham M E 2013 Identifying key controls on the behavior of an acidic-U^{VI} plume in the Savannah River Site using reactive transport modeling *J. Contam. Hydrol.* **151** 34–54
- Begg J D, Edelman C, Zavarin M and Kersting A B 2018 Sorption kinetics of plutonium(V)/(VI) to three montmorillonite clays *Appl. Geochem.* **96** 131–7
- Begg J D, Zavarin M and Kersting A B 2017 Desorption of plutonium from montmorillonite: an experimental and modeling study *Geochim. Cosmochim. Acta* **197** 278–93
- Beller H R 2005 Anaerobic, nitrate-dependent oxidation of U^{IV} oxide minerals by the chemolithoautotrophic bacterium *thiobacillus denitrificans* *Appl. Environ. Microbiol.* **71** 2170–4
- Beller H R et al 2014 Divergent aquifer biogeochemical systems converge on similar and unexpected Cr^{VI} reduction products *Environ. Sci. Technol.* **48** 10699–706
- Bender J, Duff M, Phillips P and Hill M 2000 Bioremediation and bioreduction of dissolved U^{VI} by microbial mat consortium supported on silica gel particles *Environ. Sci. Technol.* **34** 3235–41
- Berkowitz B 2002 Characterizing flow and transport in fractured geological media: a review *Adv. Water Resour.* **25** 861–84
- Bernhard G, Geipel G, Reich T, Brendler V, Amayri S and Nitsche H 2001 Uranyl(VI) carbonate complex formation: validation of the Ca₂UO₂(CO₃)_{3aq} species *Radiochim. Acta* **89** 511–18
- Bernhardt E S, Blaszcak J R, Ficken C D, Fork M L, Kaiser K E and Seybold E C 2017 Control points in ecosystems: moving beyond the hot spot hot moment concept *Ecosystems* **20** 665–82

- Bi Y, Stylo M, Bernier-Latmani R and Hayes K F 2016 Rapid mobilization of noncrystalline U^{IV} coupled with FeS oxidation *Environ. Sci. Technol.* **50** 1403–11
- Binley A, Hubbard S S, Huisman J A, Revil A, Robinson D A, Singha K and Slater L D 2015 The emergence of hydrogeophysics for improved understanding of subsurface processes over multiple scales *Water Resour. Res.* **51** 3837–66
- Blute N K, Brabander D J, Hemond H F, Sutton S R, Newville M G and Rivers M L 2004 Arsenic sequestration by ferric iron plaque on cattail roots *Environ. Sci. Technol.* **38** 6074–7
- Bond D L and Fendorf S 2003 Kinetics and structural constraints of chromate reduction by green rusts *Environ. Sci. Technol.* **37** 2750–7
- Bone S E, Cahill M R, Jones M E, Fendorf S, Davis J, Williams K H and Bargar J R 2017a Oxidative uranium release from anoxic sediments under diffusion-limited conditions *Environ. Sci. Technol.* **51** 11039–47
- Bone S E, Cliff J, Weaver K, Takacs C J, Roycroft S, Fendorf S and Bargar J R 2020 Complexation by organic matter controls uranium mobility in anoxic sediments *Environ. Sci. Technol.* **54** 1493–502
- Bone S E, Dynes J J, Cliff J and Bargar J R 2017b U^{IV} adsorption by natural organic matter in anoxic sediments *Proc. Natl Acad. Sci.* **114** 711–6
- Boyanov M I, Fletcher K E, Kwon M J, Rui X, O'Loughlin E J, Löffler F E and Kemner K M 2011 Solution and microbial controls on the formation of reduced U^{IV} species *Environ. Sci. Technol.* **45** 8336–44
- Boyanov M I, Latta D E, Scherer M M, O'Loughlin E J and Kemner K M 2017b Surface area effects on the reduction of U^{VI} in the presence of synthetic montmorillonite *Chem. Geol.* **464** 110–17
- Boyanov M I, O'Loughlin E J, Roden E E, Fein J B and Kemner K M 2007 Adsorption of Fe^{II} and U^{VI} to carboxyl-functionalized microspheres: the influence of speciation on uranyl reduction studied by titration and XAFS *Geochim. Cosmochim. Acta* **71** 1898–1912
- Boyanov M, Kelly S, Kemner K, Bunker B, Fein J and Fowle D 2003 Adsorption of cadmium to *Bacillus subtilis* bacterial cell walls: a pH-dependent X-ray absorption fine structure spectroscopy study *Geochim. Cosmochim. Acta* **67** 3299–3311
- Boyanov M, Latta D E, Mishra B, Scherer M M, O'Loughlin E J and Kemner K 2016 The effect of iron oxides and montmorillonite clays on the trans-formations of uranium under reducing conditions *3rd Int. Workshop on Advanced Techniques in Actinide Spectroscopy* (Richland, WA)
- Boyanov M, Latta D E, Mishra B, Scherer M M, O'Loughlin E J and Kemner K 2017a Stabilization of a mixed-valence U^V - U^{VI} phase in systems with reduced SWy-2 and NAU-1 clays *Goldschmidt (Paris, France, 15 August)*
- Boye K, Noël V, Tfaily M M, Bone S E, Williams K H, Bargar J R and Fendorf S 2017 Thermodynamically controlled preservation of organic carbon in floodplains *Nat. Geosci.* **10** 415–19
- Brantley S L et al 2017 Designing a network of critical zone observatories to explore the living skin of the terrestrial Earth *Earth Surf. Dyn.* **5** 841–60
- Brender J D et al 2013 Prenatal nitrate intake from drinking water and selected birth defects in offspring of participants in the national birth defects prevention study *Environ. Health Perspect.* **121** 1083–9
- Brooks S C, Fredrickson J K, Carroll S L, Kennedy D W, Zachara J M, Plymale A E, Kelly S D, Kemner K M and Fendorf S 2003 Inhibition of bacterial U^{VI} reduction by calcium *Environ. Sci. Technol.* **37** 1850–8
- Brooks S C, Riscassi A L and Miller C L 2021 Diel mercury concentration variations in a mercury-impacted stream (in preparation)
- Brooks S C and Southworth G R 2011 History of mercury use and environmental contamination at the Oak Ridge Y-12 Plant *Environ. Pollut.* **159** 219–28
- Brown C T et al 2015 Unusual biology across a group comprising more than 15% of domain bacteria *Nature* **523** 208–11
- Brown L R and Halweil B 1998 China's water shortage could shake world food security *World Watch* **11** 10–16
- Campbell K M et al 2012 Geochemical, mineralogical and microbiological characteristics of sediment from a naturally reduced zone in a uranium-contaminated aquifer *Appl. Geochem.* **27** 1499–1511
- Cardarelli E L, Bargar J R and Francis C A 2020 Diverse Thaumarchaeota dominate subsurface ammonia-oxidizing communities in semi-arid floodplains in the Western United States *Microbial Ecol.* **80** 778–92
- Carnevali P B M et al 2021 Meanders as a scaling motif for understanding of floodplain soil microbiome and biogeochemical potential at the watershed scale *Microbiome* **9** 1–23
- Carrell A A et al 2021 Nutrient exposure alters microbial composition, structure and mercury methylating activity in periphyton in a contaminated watershed *Front. Microbiol.* **12** 543
- Castelle C J et al 2013 Extraordinary phylogenetic diversity and metabolic versatility in aquifer sediment *Nat. Commun.* **4** 1–10
- Cerrato J M, Ashner M N, Alessi D S, Lezama-Pacheco J S, Bernier-Latmani R, Bargar J R and Giammar D E 2013 Relative reactivity of biogenic and chemogenic uraninite and biogenic noncrystalline U^{IV} *Environ. Sci. Technol.* **47** 9756–63
- Chadwick K D et al 2020a NEON AOP foliar trait maps, maps of model uncertainty estimates and conifer map, East River, CO 2018 ESS-DIVE repository (<https://doi.org/10.15485/1618133>)
- Chadwick K D et al 2020b Locations, metadata and species cover from field sampling survey associated with NEON AOP survey, East River, CO 2018 ESS-DIVE repository (<https://doi.org/10.15485/1618130>)
- Chadwick K D, Grant K, Henderson A, Scott A, McCormick M, Pierce S, Hastings Porro M and Maher K 2020c Leaf mass per area and leaf water content measurements from field survey in association with NEON AOP survey, East River, CO 2018 ESS-DIVE repository (<https://doi.org/10.15485/1618132>)
- Chang H-s, Buettner S W, Seaman J C, Jaffé P R, Koster van Groos P G, Li D, Peacock A D, Scheckel K G and Kaplan D I 2014 Uranium immobilization in an iron-rich rhizosphere of a native wetland plant from the Savannah River Site under reducing conditions *Environ. Sci. Technol.* **48** 9270–8
- Chen H, Johnston R C, Mann B F, Chu R K, Tolic N, Parks J M and Gu B 2017 Identification of mercury and dissolved organic matter complexes using ultrahigh resolution mass spectrometry *Environ. Sci. Technol. Lett.* **4** 59–65
- Chen J, Hubbard S, Rubin Y, Murray C, Roden E and Majer E 2004 Geochemical characterization using geophysical data and Markov Chain Monte Carlo methods: a case study at the South Oyster bacterial transport site in Virginia *Water Resour. Res.* **40** W12412
- Chen X et al 2021 Integrating field observations and process-based modeling to predict watershed water quality under environmental perturbations *J. Hydrol.* **602** 125762
- Christensen N S and Lettenmaier D P 2007 A multimodel ensemble approach to assessment of climate change impacts on the hydrology and water resources of the Colorado River Basin *Hydrol. Earth Syst. Sci.* **11** 1417–34
- Colombo R, Meroni M, Marchesi A, Busetto L, Rossini M, Giardino C and Panigada C 2008 Estimation of leaf and canopy water content in poplar plantations by means of hyperspectral indices and inverse modeling *Remote Sens. Environ.* **112** 1820–34
- Coon E T, Moulton J D, Kikinzon E, Berndt M, Manzini G, Garimella R, Lipnikov K and Painter S L 2020 Coupling surface flow and subsurface flow in complex soil structures

- using mimetic finite differences *Adv. Water Resour.* **144** 103701
- Coon E T, Moulton J D and Painter S L 2016 Managing complexity in simulations of land surface and near-surface processes *Environ. Model. Softw.* **78** 134–49
- Cumberland S A, Douglas G, Grice K and Moreau J W 2016 Uranium mobility in organic matter-rich sediments: a review of geological and geochemical processes *Earth-Sci. Rev.* **159** 160–85
- Curtis G P, Davis J A and Naftz D L 2006 Simulation of reactive transport of uranium(VI) in groundwater with variable chemical conditions *Water Resour. Res.* **42** W04404
- Curtis G P, Fox P, Kohler M and Davis J A 2004 Comparison of *in situ* uranium K_D values with a laboratory determined surface complexation model *Appl. Geochem.* **19** 1643–53
- Dafflon B, Irving J and Barrash W 2011 Inversion of multiple intersecting high-resolution crosshole GPR profiles for hydrological characterization at the Boise Hydrogeophysical Research Site *J. Appl. Geophys.* **73** 305–14
- Dafflon B, Malenda H, and Dwivedi D 2020 Groundwater level elevation and temperature across Meander C at the Lower Montane in the East River Watershed, Colorado *Technical Report* Environmental System Science Data Infrastructure for a Virtual Ecosystem.
- Dai M, Kelley J M and Buesseler K O 2002 Sources and migration of plutonium in groundwater at the Savannah River Site *Environ. Sci. Technol.* **36** 3690–9
- Dam W L, Campbell S, Johnson R H, Looney B B, Denham M E, Eddy-Dilek C A and Babits S J 2015 Refining the site conceptual model at a former uranium mill site in Riverton, Wyoming, USA *Environ. Earth Sci.* **74** 7255–65
- Dangelmayr M A, Reimus P W, Wasserman N L, Punsal J J, Johnson R H, Clay J T and Stone J J 2017 Laboratory column experiments and transport modeling to evaluate retardation of uranium in an aquifer downgradient of a uranium *in-situ* recovery site *Appl. Geochem.* **80** 1–13
- Davis J A, Meece D E, Kohler M and Curtis G P 2004 Approaches to surface complexation modeling of uranium(VI) adsorption on aquifer sediments *Geochim. Cosmochim. Acta* **68** 3621–41
- Davis J, Coston J, Kent D and Fuller C 1998 Application of the surface complexation concept to complex mineral assemblages *Environ. Sci. Technol.* **32** 2820–8
- Deblonde G J-P, Kersting A B and Zavarin M 2020 Open questions on the environmental chemistry of radionuclides *Commun. Chem.* **3** 1–5
- DeFlaun M F, Fuller M E, Zhang P, Johnson W P, Mailloux B J, Holben W E, Kovacic W P, Balkwill D L and Onstott T C 2001 Comparison of methods for monitoring bacterial transport in the subsurface *J. Microbiol. Methods* **47** 219–31
- Demers J D, Blum J D, Brooks S C, Donovan P M, Riscassi A L, Miller C L, Zheng W and Gu B 2018 Hg isotopes reveal in-stream processing and legacy inputs in East Fork Poplar Creek, Oak Ridge, Tennessee, USA *Environ. Sci.: Process. Impacts* **20** 686–707
- Dempsey D, Simmons S, Archer R and Rowland J 2012 Delineation of catchment zones of geothermal systems in large-scale rifted settings *J. Geophys. Res.: Solid Earth* **117** B10201
- Devadoss J, Falco N, Dafflon B, Wu Y, Franklin M, Hermes A, Hinckley E-L S and Wainwright H 2020 Remote sensing-informed zonation for understanding snow, plant and soil moisture dynamics within a mountain ecosystem *Remote Sens.* **12** 2733
- Dewey C, Sokaras D, Kroll T, Bargar J R and Fendorf S 2020 Calcium-uranyl-carbonate species kinetically limit U^{VI} reduction by Fe^{II} and lead to U^V -bearing ferrihydrite *Environ. Sci. Technol.* **54** 6021–30
- Diamond S et al 2019 Mediterranean grassland soil C–N compound turnover is dependent on rainfall and depth and is mediated by genomically divergent microorganisms *Nat. Microbiol.* **4** 1356–67
- Dodds W K 2006 Nutrients and the “dead zone”: the link between nutrient ratios and dissolved oxygen in the northern Gulf of Mexico *Front. Ecol. Environ.* **4** 211–17
- DOE-Complexity 2009 Complex systems science for subsurface fate and transport (available at: https://doesbr.org/complexityreport/SubsurfaceComplexitywithCover_03-05-10HR200dpi.pdf) (Accessed 15 March 2021)
- DOE-LM 1998 Final site observational work plan for the UMTRA project site at Riverton, Wyoming *Document U0013801* Grand Junction, CO (available at: www.lm.doe.gov/Riverton/RVT00000133.pdf) (Accessed 15 March 2021)
- DOE, Nevada Operations Office 2000 *United States Nuclear Tests: July 1945 Through September 1992* (Oak Ridge, TN: Department of Energy, Nevada Operations Office) (available at: www.osti.gov/scitech)
- DOE 1996 *The 1996 Baseline Environmental Management Report (Volume I and II)* (Washington, DC: DOE/EM-0319, Office of Environmental Management)
- DOE 1997 *Linking Legacies: Connecting the Cold War Nuclear Weapons Processes to Their Environmental Consequences* (Washington, DC: DOE/EM-0319, US DOE Office of Environmental Management)
- Dong H, Onstott T C, DeFlaun M F, Fuller M E, Scheibe T D, Streger S H, Rothmel R K and Mailloux B J 2002a Relative dominance of physical versus chemical effects on the transport of adhesion-deficient bacteria in intact cores from South Oyster, Virginia *Environ. Sci. Technol.* **36** 891–900
- Dong H, Onstott T C, Ko C-H, Hollingsworth A D, Brown D G and Mailloux B J 2002b Theoretical prediction of collision efficiency between adhesion-deficient bacteria and sediment grain surface *Colloids Surf. B* **24** 229–45
- Dong Y et al 2020 Controls on iron reduction and biomineralization over broad environmental conditions as suggested by the Firmicutes *Orenia metallireducens* strain Z6 *Environ. Sci. Technol.* **54** 10128–40
- Donovan P M, Blum J D, Demers J D, Gu B, Brooks S C and Peryam J 2014 Identification of multiple mercury sources to stream sediments near Oak Ridge, TN, USA *Environ. Sci. Technol.* **48** 3666–74
- Droz B, Dumas N, Duckworth O W and Peña J 2015 A comparison of the sorption reactivity of bacteriogenic and mycogenic Mn oxide nanoparticles *Environ. Sci. Technol.* **49** 4200–8
- Duckworth O W, Akafia M M, Andrews M Y and Bargar J R 2014 Siderophore-promoted dissolution of chromium from hydroxide minerals *Environ. Sci.: Process. Impacts* **16** 1348–59
- Duckworth O W, Bargar J R, Jarzecki A A, Oyerinde O, Spiro T G and Sposito G 2009 The exceptionally stable cobalt(III)–desferrioxamine B complex *Marine Chem.* **113** 114–22
- Dullies F, Lutze W, Gong W and Nuttall H E 2010 Biological reduction of uranium—from the laboratory to the field *Sci. Total Environ.* **408** 6260–71
- Duncan J M, Groffman P M and Band L E 2013 Towards closing the watershed nitrogen budget: spatial and temporal scaling of denitrification *J. Geophys. Res.: Biogeosci.* **118** 1105–19
- Durrant C B, Begg J D, Kersting A B and Zavarin M 2018 Cesium sorption reversibility and kinetics on illite, montmorillonite and kaolinite *Sci. Total Environ.* **610** 511–20
- Dwivedi D et al 2018b Geochemical exports to river from the intrameander hyporheic zone under transient hydrologic conditions: East River Mountainous Watershed, Colorado *Water Resour. Res.* **54** 8456–77
- Dwivedi D, Arora B, Steefel C I, Dafflon B and Versteeg R 2018a Hot spots and hot moments of nitrogen in a riparian corridor *Water Resour. Res.* **54** 205–22
- Dwivedi D, Mital U, Faybishenko B, Dafflon B, Varadharajan C, Agarwal D, Williams K H, Steefel C I and Hubbard S S 2022 Imputation of contiguous gaps and extremes of subhourly

- groundwater time series using random forests *J. Mach. Learn. Model. Comput.* **3** 1–22
- Dwivedi D, Mohanty B P and Lesikar B J 2013 Estimating *Escherichia coli* loads in streams based on various physical, chemical and biological factors *Water Resour. Res.* **49** 2896–906
- Dwivedi D, Mohanty B P and Lesikar B J 2016b Impact of the linked surface water-soil water-groundwater system on transport of *E. coli* in the subsurface *Water Air Soil Pollut.* **227** 1–16
- Dwivedi D, Riley W, Torn M, Spycher N, Maggi F and Tang J 2017a Mineral properties, microbes, transport and plant-input profiles control vertical distribution and age of soil carbon stocks *Soil Biol. Biochem.* **107** 244–59
- Dwivedi D, Steefel C, Siirila-Woodburn E, Kallemov B, Moulton J D, Kikinzon C E, Hammond G, Foster L and Maxwell R 2016a Testing code interoperability and productivity on modeling integrated surface subsurface water flow and biogeochemical cycling in the hyporheic zone—IDEAS use case 1 *Technical Report* Annual joint investigators meeting of the Department of Energy's Office of Biological and Environmental Research (BER) at: Potomac, Maryland
- Dwivedi D, Steefel I C, Arora B and Bisht G 2017b Impact of intra-meander hyporheic flow on nitrogen cycling *Proc. Earth Planet. Sci.* **17** 404–7
- Dwivedi D, Tang J, Bouskill N, Georgiou K, Chacon S S and Riley W J 2019 Abiotic and biotic controls on soil organo-mineral interactions: developing model structures to analyze why soil organic matter persists *Rev. Mineral. Geochem.* **85** 329–48
- Dzombak D A and Morel F M 1990 *Surface Complexation Modeling: Hydrous Ferric Oxide* (New York: Wiley)
- Ellis P A, Mackay R and Rivett M O 2007 Quantifying urban river-aquifer fluid exchange processes: a multi-scale problem *J. Contam. Hydrol.* **91** 58–80
- Engel M, Boye K, Noël V, Babey T, Bargar J R and Fendorf S 2021 Simulated aquifer heterogeneity leads to enhanced attenuation and multiple retention processes of zinc *Environ. Sci. Technol.* **55** 2939–48
- Estes E, Andeer P, Nordlund D, Wankel S and Hansel C 2017 Biogenic manganese oxides as reservoirs of organic carbon and proteins in terrestrial and marine environments *Geobiology* **15** 158–72
- Falco N et al 2018 Watershed functioning zonation: advanced watershed characterization across scales *AGU Fall Meeting Abstracts* vol 2018 p H51O–1512
- Famiglietti J S 2014 The global groundwater crisis *Nat. Clim. Change* **4** 945–8
- Fan D, Anitori R P, Tebo B M, Tratnyek P G, Lezama Pacheco J S, Kukkadapu R K, Kovarik L, Engelhard M H and Bowden M E 2014 Oxidative remobilization of technetium sequestered by sulfide-transformed nano zerovalent iron *Environ. Sci. Technol.* **48** 7409–17
- Fang Y et al 2020 High-performance simulation of dynamic hydrologic exchange and implications for surrogate flow and reactive transport modeling in a large river corridor *Front. Water* **2** 564211
- Faybishenko B et al 2008 *In situ* long-term reductive bioimmobilization of Cr^{VI} in groundwater using hydrogen release compound *Environ. Sci. Technol.* **42** 8478–85
- Faybishenko B, Versteeg R, Pastorello G, Dwivedi D, Varadharajan C and Agarwal D 2021 Challenging problems of quality assurance and quality control (QA/QC) of meteorological time series data *Stochastic Environ. Res. Risk Assess.* **1436**–3259
- Feilhauer H, Asner G P and Martin R E 2015 Multi-method ensemble selection of spectral bands related to leaf biochemistry *Remote Sens. Environ.* **164** 57–65
- Felmy A R, Moore D A, Rosso K M, Qafoku O, Rai D, Buck E C and Ilton E S 2011 Heterogeneous reduction of PuO₂ with Fe^{II}: importance of the Fe^{III} reaction product *Environ. Sci. Technol.* **45** 3952–8
- Felmy A R and Rustad J R 1998 Molecular statics calculations of proton binding to goethite surfaces: thermodynamic modeling of the surface charging and protonation of goethite in aqueous solution *Geochim. Cosmochim. Acta* **62** 25–31
- Fendorf S E and Li G 1996 Kinetics of chromate reduction by ferrous iron *Environ. Sci. Technol.* **30** 1614–17
- Finneran K T, Housewright M E and Lovley D R 2002 Multiple influences of nitrate on uranium solubility during bioremediation of uranium-contaminated subsurface sediments *Environ. Microbiol.* **4** 510–16
- Finsterle S, Sonnenthal E L and Spycher N 2014 Advances in subsurface modeling using the TOUGH suite of simulators *Comput. Geosci.* **65** 2–12
- Fletcher K E, Boyanov M I, Thomas S H, Wu Q, Kemner K M and Löffler F E 2010 U(VI) reduction to mononuclear U^{IV} by *Desulfitobacterium* species *Environ. Sci. Technol.* **44** 4705–9
- Flores Orozco A, Bucker M and Williams K 2014 Characterization of natural attenuation in a uranium-contaminated site by means of induced polarization imaging *EGU General Conf. Abstracts* p 8942
- Flores Orozco A, Gallistl J, Bucker M and Williams K H 2018 Decay curve analysis for data error quantification in time-domain induced polarization imaging *Geophysics* **83** E75–E86
- Flores Orozco A, Williams K H, Long P E, Hubbard S S and Kemna A 2011 Using complex resistivity imaging to infer biogeochemical processes associated with bioremediation of an uranium-contaminated aquifer *J. Geophys. Res.: Biogeosci.* **116** G03001
- Flynn T M, O'Loughlin E J, Mishra B, DiChristina T J and Kemner K M 2014 Sulfur-mediated electron shuttling during bacterial iron reduction *Science* **344** 1039–42
- Fogg G E, Noyes C D and Carle S F 1998 Geologically based model of heterogeneous hydraulic conductivity in an alluvial setting *Hydrogeol. J.* **6** 131–43
- Fox P M, Davis J A, Hay M B, Conrad M E, Campbell K M, Williams K H and Long P E 2012 Rate-limited U^{VI} desorption during a small-scale tracer test in a heterogeneous uranium-contaminated aquifer *Water Resour. Res.* **48** W05512
- Fox P M, Davis J A and Zachara J M 2006 The effect of calcium on aqueous uranium(VI) speciation and adsorption to ferrihydrite and quartz *Geochim. Cosmochim. Acta* **70** 1379–87
- Frazier S W, Kretzschmar R and Kraemer S M 2005 Bacterial siderophores promote dissolution of UO₂ under reducing conditions *Environ. Sci. Technol.* **39** 5709–15
- Fredrickson J K, Zachara J M, Kennedy D W, Dong H, Onstott T C, Hinman N W and Li S-m 1998 Biogenic iron mineralization accompanying the dissimilatory reduction of hydrous ferric oxide by a groundwater bacterium *Geochim. Cosmochim. Acta* **62** 3239–57
- Fredrickson J K, Zachara J M, Kennedy D W, Duff M C, Gorby Y A, Shu-mei W L and Krupka K M 2000 Reduction of U^{VI} in goethite (α -FeOOH) suspensions by a dissimilatory metal-reducing bacterium *Geochim. Cosmochim. Acta* **64** 3085–98
- Fredrickson J K, Zachara J M, Kennedy D W, Kukkadapu R K, McKinley J P, Heald S M, Liu C and Plymale A E 2004 Reduction of TcO₄⁻ by sediment-associated biogenic Fe^{III} *Geochim. Cosmochim. Acta* **68** 3171–87
- Fritz B G and Arntzen E V 2007 Effect of rapidly changing river stage on uranium flux through the hyporheic zone *Groundwater* **45** 753–60
- Fuller C C and Bargar J R 2014 Processes of zinc attenuation by biogenic manganese oxides forming in the hyporheic zone of Pinal Creek, Arizona *Environ. Sci. Technol.* **48** 2165–72
- Fuller M E, Streger S H, Rothmel R K, Mailloux B J, Hall J A, Onstott T C, Fredrickson J K, Balkwill D L and DeFlaun M F 2000 Development of a vital fluorescent staining method for

- monitoring bacterial transport in subsurface environments *Appl. Environ. Microbiol.* **66** 4486–96
- Garimella R V, Perkins W A, Buksas M W, Berndt M, Lipnikov K, Coon E, Moulton J D and Painter S L 2014 Mesh infrastructure for coupled multiprocess geophysical simulations *Proc. Eng.* **82** 34–45
- Gentine P, Pritchard M, Rasp S, Reinaudi G and Yacalis G 2018 Could machine learning break the convection parameterization deadlock? *Geophys. Res. Lett.* **45** 5742–51
- Geszvain K et al 2012 The molecular biogeochemistry of manganese(II) oxidation *Biochem. Soc. Trans.* **40** 1244–8
- Ghiorse W and Wobber F 1989 Special issue on deep subsurface microbiology-introduction
- Gihring T M et al 2011 A limited microbial consortium is responsible for extended bioreduction of uranium in a contaminated aquifer *Appl. Environ. Microbiol.* **77** 5955–65
- Gillin C P, Bailey S W, McGuire K J and Gannon J P 2015 Mapping of hydrogeologic spatial patterns in a steep headwater catchment *Soil Sci. Soc. Am. J.* **79** 440–53
- Gilmour C C et al 2013 Mercury methylation by novel microorganisms from new environments *Environ. Sci. Technol.* **47** 11810–20
- Gilson E R, Huang S, Koster van Groos P G, Scheckel K G, Qafoku O, Peacock A D, Kaplan D I and Jaffé P R 2015 Uranium redistribution due to water table fluctuations in sandy wetland mesocosms *Environ. Sci. Technol.* **49** 12214–22
- Ginder-Vogel M, Criddle C S and Fendorf S 2006 Thermodynamic constraints on the oxidation of biogenic UO_2 by Fe^{III} (hydr) oxides *Environ. Sci. Technol.* **40** 3544–50
- Gomez-Velez J D and Harvey J W 2014 A hydrogeomorphic river network model predicts where and why hyporheic exchange is important in large basins *Geophys. Res. Lett.* **41** 6403–12
- Gorby Y A et al 2006 Electrically conductive bacterial nanowires produced by *Shewanella oneidensis* strain MR-1 and other microorganisms *Proc. Natl Acad. Sci.* **103** 11358–63
- Gorby Y A and Lovley D R 1992 Enzymic uranium precipitation *Environ. Sci. Technol.* **26** 205–7
- Greskowiak J, Prommer H, Liu C, Post V, Ma R, Zheng C and Zachara J M 2010 Comparison of parameter sensitivities between a laboratory and field-scale model of uranium transport in a dual domain, distributed rate reactive system *Water Resour. Res.* **46** W09509
- Gross T 2017 The world's threatened rivers - in pictures (available at: www.npr.org/2017/04/13/523717152/how-can-the-colorado-river-continue-to-support-36-million-people-in-7-states) (Accessed 15 March 2021)
- Gu B, Mishra B, Miller C, Wang W, Lai B, Brooks S C, Kemner K M and Liang L 2014 X-ray fluorescence mapping of mercury on suspended mineral particles and diatoms in a contaminated freshwater system *Biogeochemistry* **11** 5259–67
- Gu B, Schmitt J, Chen Z, Liang L and McCarthy J F 1994 Adsorption and desorption of natural organic matter on iron oxide: mechanisms and models *Environ. Sci. Technol.* **28** 38–46
- Gu B, Schmitt J, Chen Z, Liang L and McCarthy J F 1995 Adsorption and desorption of different organic matter fractions on iron oxide *Geochim. Cosmochim. Acta* **59** 219–29
- Gu B, Yan H, Zhou P, Watson D B, Park M and Istok J 2005 Natural humics impact uranium bioreduction and oxidation *Environ. Sci. Technol.* **39** 5268–75
- Hammond G E and Lichtner P C 2010 Field-scale model for the natural attenuation of uranium at the Hanford 300 Area using high-performance computing *Water Resour. Res.* **46** W09527
- Hammond G E, Lichtner P C and Mills R 2014 Evaluating the performance of parallel subsurface simulators: an illustrative example with PFLOTRAN *Water Resour. Res.* **50** 208–28
- Hammond G, Lichtner P and Lu C 2007 Subsurface multiphase flow and multicomponent reactive transport modeling using high-performance computing *Journal of Physics: Conf. Series* vol 78 (IOP Publishing) p 012025
- Hammond G, Lichtner P, Lu C and Mills R T eds. 2012 PFLOTRAN: Reactive flow & transport code for use on laptops to leadership-class supercomputers *Groundwater Reactive Transp. Models* (Sharjah: Bentham Science Publishers) Zhang F, Gour-Tsyh (George) Y and Parker J C. 141–59
- Hansel C M, Lentini C J, Tang Y, Johnston D T, Wankel S D and Jardine P M 2015 Dominance of sulfur-fueled iron oxide reduction in low-sulfate freshwater sediments *ISME J.* **9** 2400–12
- Hansel C M, Zeiner C A, Santelli C M and Webb S M 2012 Mn(II) oxidation by an ascomycete fungus is linked to superoxide production during asexual reproduction *Proc. Natl Acad. Sci.* **109** 12621–5
- Hao Y, Sun Y and Nitao J 2012 Overview of NUFT: A versatile numerical model for simulating flow and reactive transport in porous media *Groundwater Reactive Transp. Models* (Oak Park, IL: Bentham Science Publishers) Zhang Fan, Gour-Tsyh (George) Yeh and Parker Jack C. 212–39
- Hausladen D M and Fendorf S 2017 Hexavalent chromium generation within naturally structured soils and sediments *Environ. Sci. Technol.* **51** 2058–67
- Hazen T C and Tabak H H 2005 Developments in bioremediation of soils and sediments polluted with metals and radionuclides: 2. Field research on bioremediation of metals and radionuclides *Rev. Environ. Sci. Bio/Technol.* **4** 157–83
- He C, Keren R, Whittaker M L, Farag I F, Doudna J A, Cate J H and Banfield J F 2021 Genome-resolved metagenomics reveals site-specific diversity of epibiotic CPR bacteria and DPANN archaea in groundwater ecosystems *Nat. Microbiol.* **6** 354–65
- Heathwaite A 2010 Multiple stressors on water availability at global to catchment scales: understanding human impact on nutrient cycles to protect water quality and water availability in the long term *Freshwater Biol.* **55** 241–57
- Howarth R W et al 1992 *Sulphur Cycling on the Continents: Wetlands, Terrestrial Ecosystems and Associated Water Bodies* (Chichester: John Wiley Sons Ltd)
- Hsi C-k D and Langmuir D 1985 Adsorption of uranyl onto ferric oxyhydroxides: application of the surface complexation site-binding model *Geochim. Cosmochim. Acta* **49** 1931–41
- Hubbard S S et al 2013 Quantifying and relating land-surface and subsurface variability in permafrost environments using LiDAR and surface geophysical datasets *Hydrogeol. J.* **21** 149–69
- Hubbard S S et al 2018 The East River, Colorado, Watershed: a mountainous community tested for improving predictive understanding of multiscale hydrological–biogeochemical dynamics *Vadose Zone J.* **17** 1–25
- Hubbard S S, Chen J, Peterson J, Majer E L, Williams K H, Swift D J, Mailloux B and Rubin Y 2001 Hydrogeological characterization of the South Oyster Bacterial Transport Site using geophysical data *Water Resour. Res.* **37** 2431–56
- Hubbard S S and Rubin Y 2005 Introduction to hydrogeophysics *Hydrogeophysics* (Berlin: Springer) pp 3–21
- Hubbard S S, Varadharajan C, Wu Y, Wainwright H and Dwivedi D 2020 Emerging technologies and radical collaboration to advance predictive understanding of watershed hydrobiogeochemistry *Hydrol. Process.* **34** 3175–82
- Hug L A et al 2016 A new view of the tree of life *Nat. Microbiol.* **1** 1–6
- Hug L A, Castelle C J, Wrighton K C, Thomas B C, Sharon I, Frischkorn K R, Williams K H, Tringe S G and Banfield J F 2013 Community genomic analyses constrain the distribution of metabolic traits across the Chloroflexi phylum and indicate roles in sediment carbon cycling *Microbiome* **1** 1–17
- Hug L A, Thomas B C, Brown C T, Frischkorn K R, Williams K H, Tringe S G and Banfield J F 2015 Aquifer environment

- selects for microbial species cohorts in sediment and groundwater *ISME Journal* **9** 1846–56
- Ilton E S, Haiduc A, Cahill C L and Felmy A R 2005 Mica surfaces stabilize pentavalent uranium *Inorg. Chem.* **44** 2986–8
- Islam M, Islam F and IWA W 2010 IWA Water Wiki (available at: www.iwapublishing.com/news/arsenic-contamination-groundwater-bangladesh-environmental-and-social-disaster#:~:text=The%20contamination%20of%20groundwater%20by,F%2C%202002.) (Accessed 11 March 2021)
- Istok J, Senko J, Krumholz L R, Watson D, Bogle M A, Peacock A, Chang Y-J and White D C 2004 *In situ* bioreduction of technetium and uranium in a nitrate-contaminated aquifer *Environ. Sci. Technol.* **38** 468–75
- Ithurbide A, Peulon S, Miserque F, Beaucaire C and Chaussé A 2009 Interaction between uranium(VI) and siderite (FeCO₃) surfaces in carbonate solutions *Radiochim. Acta* **97** 177–80
- Jacques D and Šimůnek J 2005 User manual of the multicomponent variably-saturated flow and transport model HP1 *Technical Report* SCK-CEN
- Jacques D, Šimůnek J, Mallants D and Van Genuchten M T 2006 Operator-splitting errors in coupled reactive transport codes for transient variably saturated flow and contaminant transport in layered soil profiles *J. Contam. Hydrol.* **88** 197–218
- Jacques D, Šimůnek J, Mallants D, van Genuchten M T and Kodešová R 2013 The HPx reactive transport models: summary of recent developments and applications *Proc. of the 4th Int. Conf., HYDRUS Software Applications to Subsurface Flow and Contaminant Transport Problems, Invited Paper, 21–22 March, 2013* (Dep. of Soil Science and Geology, Czech Univ. of Life Sciences Prague, Czech) pp 7–16
- Jacques D, Smith C, Šimůnek J and Smiles D 2012 Inverse optimization of hydraulic, solute transport and cation exchange parameters using HP1 and UCODE to simulate cation exchange *J. Contam. Hydrol.* **142** 109–25
- Jan A, Coon E T and Painter S L 2021 Toward more mechanistic representations of biogeochemical processes in river networks: implementation and demonstration of a multiscale model *Environ. Model. Softw.* **145** 105166
- Janot N et al 2016 Physico-chemical heterogeneity of organic-rich sediments in the Rifle aquifer, CO: Impact on uranium biogeochemistry *Environ. Sci. Technol.* **50** 46–53
- Jardine P, Fendorf S, Mayes M, Larsen I, Brooks S and Bailey W 1999 Fate and transport of hexavalent chromium in undisturbed heterogeneous soil *Environ. Sci. Technol.* **33** 2939–44
- Jemison N E, Bizjack M T, Johnson T M and Druhan J L 2020 Influence of physical and chemical hydrology on bioremediation of a U-contaminated aquifer informed by reactive transport modeling incorporating 238U/235U ratios *Geochim. Cosmochim. Acta* **269** 303–28
- Jencso K G, McGlynn B L, Gooseff M N, Wondzell S M, Bencala K E and Marshall L A 2009 Hydrologic connectivity between landscapes and streams: transferring reach- and plot-scale understanding to the catchment scale *Water Resour. Res.* **45** W04428
- Johnson C R, Antonopoulos D A, Boyanov M I, Flynn T M, Koval J C, Kemner K M and O'Loughlin E J 2021 Reduction of Sb^V by coupled biotic-abiotic processes under sulfidogenic conditions *Heliyon* **7** e06275
- Johnson P et al 2001 Ferrographic tracking of bacterial transport in the field at the Narrow Channel Focus Area, Oyster, VA *Environ. Sci. Technol.* **35** 182–91
- Johnson T C, Versteeg R J, Rockhold M, Slater L D, Ntarlagiannis D, Greenwood W J and Zachara J 2012 Characterization of a contaminated wellfield using 3D electrical resistivity tomography implemented with geostatistical, discontinuous boundary and known conductivity constraints *Geophysics* **77** EN85–EN96
- Johnson T C, Versteeg R J, Ward A, Day-Lewis F D and Revil A 2010 Improved hydrogeophysical characterization and monitoring through parallel modeling and inversion of time-domain resistivity and induced-polarization data *Geophysics* **75** WA27–WA41
- Johnson T, Versteeg R, Thomle J, Hammond G, Chen X and Zachara J 2015 Four-dimensional electrical conductivity monitoring of stage-driven river water intrusion: accounting for water table effects using a transient mesh boundary and conditional inversion constraints *Water Resour. Res.* **51** 6177–96
- Jones M E, LaCroix R E, Zeigler J, Ying S C, Nico P S and Keiluweit M 2020 Enzymes, manganese, or iron? Drivers of oxidative organic matter decomposition in soils *Environ. Sci. Technol.* **54** 14114–23
- Jones M E, Nico P S, Ying S, Regier T, Thieme J and Keiluweit M 2018 Manganese-driven carbon oxidation at oxic–anoxic interfaces *Environ. Sci. Technol.* **52** 12349–57
- Jordan T H et al National Research Council 2001 *Basic Research Opportunities in Earth Science* (Washington, DC: The National Academies Press)
- Jung Y, Pau G S H, Finsterle S and Doughty C 2018 TOUGH3 user's guide *Univ. California, Berkeley* (available at: https://tough.lbl.gov/assets/files/Tough3/TOUGH3_Users_Guide_v2.pdf)
- Jung Y, Pau G S H, Finsterle S and Pollyea R M 2017 TOUGH3: a new efficient version of the TOUGH suite of multiphase flow and transport simulators *Comput. Geosci.* **108** 2–7
- Kantor R S, Wrighton K C, Handley K M, Sharon I, Hug L A, Castelle C J, Thomas B C and Banfield J F 2013 Small genomes and sparse metabolisms of sediment-associated bacteria from four candidate phyla *MBio* **4** e00708–13
- Kaplan D I et al 2016 Iron mineralogy and uranium-binding environment in the rhizosphere of a wetland soil *Sci. Total Environ.* **569** 53–64
- Keiluweit M, Nico P S, Kleber M and Fendorf S 2016 Are oxygen limitations under recognized regulators of organic carbon turnover in upland soils? *Biogeochemistry* **127** 157–71
- Keiluweit M, Nico P, Harmon M E, Mao J, Pett-Ridge J and Kleber M 2015 Long-term litter decomposition controlled by manganese redox cycling *Proc. Natl Acad. Sci.* **112** E5253–60
- Kelly S D, Kemner K M, Carley J, Criddle C, Jardine P M, Marsh T L, Phillips D, Watson D and Wu W-M 2008 Speciation of uranium in sediments before and after *in situ* biostimulation *Environ. Sci. Technol.* **42** 1558–64
- Kelly S, Kemner K, Fein J, Fowle D, Boyanov M, Bunker B and Yee N 2002 X-ray absorption fine structure determination of pH-dependent U-bacterial cell wall interactions *Geochim. Cosmochim. Acta* **66** 3855–71
- Kemner K M et al 2004 Elemental and redox analysis of single bacterial cells by X-ray microbeam analysis *Science* **306** 686–7
- Keppler F, Eiden R, Niedan V, Pracht J and Schöler H 2000 Halocarbons produced by natural oxidation processes during degradation of organic matter *Nature* **403** 298–301
- Kerisit S, Felmy A R and Ilton E S 2011 Atomistic simulations of uranium incorporation into iron (hydr)-oxides *Environ. Sci. Technol.* **45** 2770–6
- Kersting A, Efurd D, Finnegan D, Rokop D, Smith D and Thompson J 1999 Migration of plutonium in ground water at the Nevada Test Site *Nature* **397** 56–9
- Keyes D E et al 2013 Multiphysics simulations: challenges and opportunities *Int. J. High Perform. Comput. Appl.* **27** 4–83
- Klingbeil R, Kleinedam S, Aspiron U, Aigner T and Teutsch G 1999 Relating lithofacies to hydrofacies: outcrop-based hydrogeological characterisation of Quaternary gravel deposits *Sediment. Geol.* **129** 299–310
- Knabe D, Guadagnini A, Riva M and Engelhardt I 2021 Uncertainty analysis and identification of key parameters controlling bacteria transport within a riverbank filtration scenario *Water Resour. Res.* **57** e2020WR027911

- Kolditz O *et al* 2012 OpenGeoSys: An open-source initiative for numerical simulation of thermo-hydro-mechanical/chemical (THM/C) processes in porous media *Environ. Earth Sci.* **67** 589–99
- Kollet S J and Maxwell R M 2008 Demonstrating fractal scaling of baseflow residence time distributions using a fully-coupled groundwater and land surface model *Geophys. Res. Lett.* **35** L07402
- Kollet S J, Maxwell R M, Woodward C S, Smith S, Vanderborght J, Vereecken H and Simmer C 2010 Proof of concept of regional scale hydrologic simulations at hydrologic resolution utilizing massively parallel computer resources *Water Resour. Res.* **46** W04201
- Kuffour B N, Engdahl N B, Woodward C S, Condon L E, Kollet S and Maxwell R M 2020 Simulating coupled surface–subsurface flows with ParFlow v3. 5.0: capabilities, applications and ongoing development of an open-source, massively parallel, integrated hydrologic model *Geosci. Model Dev.* **13** 1373–97
- Kukkadapu R K, Zachara J M, Smith S C, Fredrickson J K and Liu C 2001 Dissimilatory bacterial reduction of Al-substituted goethite in subsurface sediments *Geochim. Cosmochim. Acta* **65** 2913–24
- Kumar N, Noël V, Planer-Friedrich B, Besold J, Lezama-Pacheco J, Bargar J R, Brown Jr G E, Fendorf S and Boye K 2020a Redox heterogeneities promote thioarsenate formation and release into groundwater from low arsenic sediments *Environ. Sci. Technol.* **54** 3237–44
- Kumar R *et al* 2020b Strong hydroclimatic controls on vulnerability to subsurface nitrate contamination across Europe *Nat. Commun.* **11** 1–10
- Kurosaki H, Kaplan D I and Clark S B 2014 Impact of environmental curium on plutonium migration and isotopic signatures *Environ. Sci. Technol.* **48** 13985–91
- Kwon K D, Refson K and Sposito G 2009 On the role of Mn(IV) vacancies in the photoreductive dissolution of hexagonal birnessite *Geochim. Cosmochim. Acta* **73** 4142–50
- Kwon K D, Refson K and Sposito G 2013 Understanding the trends in transition metal sorption by vacancy sites in birnessite *Geochim. Cosmochim. Acta* **101** 222–32
- Kwon M J, Boyanov M I, Antonopoulos D A, Brulc J M, Johnston E R, Skinner K A, Kemner K M and O'Loughlin E J 2014a Effects of dissimilatory sulfate reduction on Fe^{III}(hydr)oxide reduction and microbial community development *Geochim. Cosmochim. Acta* **129** 177–90
- Kwon M J, Yang J-S, Shim M J, Boyanov M I, Kemner K M and O'Loughlin E J 2014b Acid extraction overestimates the total Fe^{II} in the presence of iron (hydr) oxide and sulfide minerals *Environ. Sci. Technol. Lett.* **1** 310–14
- Labrenz M, Druschel G K, Thomsen-Ebert T, Gilbert B, Welch S A, Kemner K M, Logan G A and Summons R E 2000 Gelsomina De Stasio, Philip L. Bond, Barry Lai, Shelly D. Kelly and Jillian F. Banfield, Sphalerite (ZnS) deposits forming in natural biofilms of sulfate-reducing bacteria *Science* **290** 1744–47
- Latta D E, Boyanov M I, Kemner K M, O'Loughlin E J and Scherer M M 2012a Abiotic reduction of uranium by Fe^{II} in soil *Appl. Geochem.* **27** 1512–24
- Latta D E, Gorski C A, Boyanov M I, O'Loughlin E J, Kemner K M and Scherer M M 2012b Influence of magnetite stoichiometry on U^{VI} reduction *Environ. Sci. Technol.* **46** 778–86
- Latta D E, Kemner K M, Mishra B and Boyanov M I 2016 Effects of calcium and phosphate on uranium(IV) oxidation: comparison between nanoparticulate uraninite and amorphous U^{IV}–phosphate *Geochim. Cosmochim. Acta* **174** 122–42
- Latta D E, Mishra B, Cook R E, Kemner K M and Boyanov M I 2014 Stable U^{IV} complexes form at high-affinity mineral surface sites *Environ. Sci. Technol.* **48** 1683–91
- Learman D, Wankel S, Webb S, Martinez N, Madden A and Hansel C M 2011 Coupled biotic–abiotic Mn(II) oxidation pathway mediates the formation and structural evolution of biogenic Mn oxides *Geochim. Cosmochim. Acta* **75** 6048–63
- Lefebvre P, Noël V, Lau K V, Jemison N E, Weaver K L, Williams K H, Bargar J R and Maher K 2019 Isotopic fingerprint of uranium accumulation and redox cycling in floodplains of the Upper Colorado River Basin *Environ. Sci. Technol.* **53** 3399–409
- Lezama-Pacheco J S *et al* 2015 Long-term *in situ* oxidation of biogenic uraninite in an alluvial aquifer: impact of dissolved oxygen and calcium *Environ. Sci. Technol.* **49** 7340–7
- Li L *et al* 2017 Expanding the role of reactive transport models in critical zone processes *Earth-Sci. Rev.* **165** 280–301
- Li L, Steefel C I, Kowalsky M B, Englert A and Hubbard S S 2010 Effects of physical and geochemical heterogeneities on mineral transformation and biomass accumulation during biostimulation experiments at Rifle, Colorado *J. Contam. Hydrol.* **112** 45–63
- Li L, Steefel C I, Williams K H, Wilkins M J and Hubbard S S 2009 Mineral transformation and biomass accumulation associated with uranium bioremediation at Rifle, Colorado *Environ. Sci. Technol.* **43** 5429–35
- Lichtner P C, Hammond G E, Lu C, Karra S, Bisht G, Andre B, Mills R, and Kumar J 2015 PFLOTTRAN user manual: a massively parallel reactive flow and transport model for describing surface and subsurface processes *Technical Report* Los Alamos National Lab. (LANL), Los Alamos, NM (United States); Sandia
- Ling F T, Post J E, Heaney P J, Santelli C M, Ilton E S, Burgos W D and Rose A W 2020 A multi-method characterization of natural terrestrial birnessites *Am. Mineral.* **105** 833–47
- Liu Y, Xu F and Liu C 2017 Coupled hydro-biogeochemical processes controlling Cr reductive immobilization in Columbia River hyporheic zone *Environ. Sci. Technol.* **51** 1508–17
- Loar J M, Stewart A J and Smith J G 2011 Twenty-five years of ecological recovery of East Fork Poplar Creek: review of environmental problems and remedial actions *Environ. Manage.* **47** 1010–20
- Long J C S, Mitchell J K, Randall J, Jeffrey J, Fischer J N, Illangasekare T H, Mills A L, Reed D T, Sacks J, Scanlon B R, Silver L T and Wely C National Research Council 2000 *Research Needs in Subsurface Science* (Washington, DC: National Academies Press)
- Long P E *et al* 2012 Microbiological, geochemical and hydrologic processes controlling uranium mobility: an integrated field scale subsurface research challenge site at Rifle, Colorado, February 2011 to January 2012 *Technical Report* Lawrence Berkeley National Lab. (LBNL), Berkeley, CA (United States)
- Lovley D R 1993 Dissimilatory metal reduction *Annu. Rev. Microbiol.* **47** 263–90
- Lovley D R 2000 *Environ. Microbe-Metal Interact.* (Washington DC: ASM Press) 1–30
- Lovley D R 2003 Cleaning up with genomics: applying molecular biology to bioremediation *Nat. Rev. Microbiol.* **1** 35–44
- Lovley D R and Phillips E J 1992 Bioremediation of uranium contamination with enzymatic uranium reduction *Environ. Sci. Technol.* **26** 2228–34
- Lovley D R, Stolz J F, Nord G L and Phillips E J 1987 Anaerobic production of magnetite by a dissimilatory iron-reducing microorganism *Nature* **330** 252–4
- Lu J, Tian Z, Yu J, Yang M and Zhang Y 2018 Distribution and abundance of antibiotic resistance genes in sand settling reservoirs and drinking water treatment plants across the Yellow River, China *Water* **10** 246
- Luef B *et al* 2015 Diverse uncultivated ultra-small bacterial cells in groundwater *Nat. Commun.* **6** 1–8
- Luo W and Gu B 2009 Dissolution and mobilization of uranium in a reduced sediment by natural humic substances under anaerobic conditions *Environ. Sci. Technol.* **43** 152–6
- Luo Y, Keenan T F and Smith M 2015 Predictability of the terrestrial carbon cycle *Global Change Biol.* **21** 1737–51
- Madritch M D, Kingdon C C, Singh A, Mock K E, Lindroth R L and Townsend P A 2014 Imaging spectroscopy links aspen

- genotype with below-ground processes at landscape scales *Phil. Trans. R. Soc. B* **369** 20130194
- Maggi F *et al* 2008 A mechanistic treatment of the dominant soil nitrogen cycling processes: model development, testing and application *J. Geophys. Res.: Biogeosci.* **113** G02016
- Maher K, Bargar J R and Brown Jr G E 2013 Environmental speciation of actinides *Inorg. Chem.* **52** 3510–32
- Mailloux B J *et al* 2003 The role of physical, chemical and microbial heterogeneity on the field-scale transport and attachment of bacteria *Water Resour. Res.* **39** 1142
- Massey M S, Lezama-Pacheco J S, Michel F M and Fendorf S 2014 Uranium incorporation into aluminum-substituted ferrihydrite during iron(II)-induced transformation *Environ. Sci.: Process. Impacts* **16** 2137–44
- Mayer K U, Frind E O and Blowes D W 2002 Multicomponent reactive transport modeling in variably saturated porous media using a generalized formulation for kinetically controlled reactions *Water Resour. Res.* **38** 13–1
- Mayes M, Jardine P, Larsen I, Brooks S and Fendorf S 2000 Multispecies transport of metal–EDTA complexes and chromate through undisturbed columns of weathered fractured saprolite *J. Contam. Hydrol.* **45** 243–65
- McCarthy J F, Czerwinski K R, Sanford W E, Jardine P M and Marsh J D 1998 Mobilization of transuranic radionuclides from disposal trenches by natural organic matter *J. Contam. Hydrol.* **30** 49–77
- McClain C N, Fendorf S, Webb S M and Maher K 2017 Quantifying Cr^{VI} production and export from serpentine soil of the California coast range *Environ. Sci. Technol.* **51** 141–9
- Megonigal J P, Hines M and Visscher P 2004 *Biogeochemistry* ed H D Holland, W H Schlesinger and K K Turekian (Oxford: Elsevier-Pergamon Oxford) **8** 317–424
- Megonigal J P and Neubauer S C 2019 *Biogeochemistry of tidal freshwater wetlands Coastal Wetlands* (Amsterdam: Elsevier) pp 641–83
- Meile C and Scheibe T D 2019 Reactive transport modeling of microbial dynamics *Elem.: Int. Magazine Mineral. Geochem. Petrol.* **15** 111–16
- Mital U, Dwivedi D, Brown J B, Faybisenko B, Painter S L and Steefel C I 2020 Sequential imputation of missing spatio-temporal precipitation data using random forests *Front. Water* **2** 20
- Moon H S, Komlos J and Jaffé P R 2007 Uranium reoxidation in previously bioreduced sediment by dissolved oxygen and nitrate *Environ. Sci. Technol.* **41** 4587–92
- Moon H, Komlos J and Jaffe P R 2009 Biogenic U^{IV} oxidation by dissolved oxygen and nitrate in sediment after prolonged U^{VI}/Fe^{III}/SO₄²⁻ reduction *J. Contam. Hydrol.* **105** 18–27
- Morin G *et al* 2016 Mononuclear U^{IV} complexes and ningyoite as major uranium species in lake sediments *Geochem. Perspect. Lett.* **2** 95–105
- Mulholland P J *et al* 2008 Stream denitrification across biomes and its response to anthropogenic nitrate loading *Nature* **452** 202–5
- N'Guessan A L, Moon H S, Peacock A D, Tan H, Sinha M, Long P E and Jaffé P R 2010 Postbiostimulation microbial community structure changes that control the reoxidation of uranium *FEMS Microbiol. Ecol.* **74** 184–95
- NABIR 2006 Introducing NABIR (available at: www2.lbl.gov/NABIRarchive/generalinfo/intro.html) (Accessed 15 March 2021)
- Nagel T, Shao H, Singh A, Watanabe N, Roßkopf C, Linder M, Wörner A and Kolditz O 2013 Non-equilibrium thermochemical heat storage in porous media: part 1—Conceptual model *Energy* **60** 254–70
- Nealson K H and Saffarini D 1994 Iron and manganese in anaerobic respiration: environmental significance, physiology and regulation *Annu. Rev. Microbiol.* **48** 311–44
- Neeway J J, Kaplan D I, Bagwell C E, Rockhold M L, Szecsody J E, Truex M J and Qafoku N P 2019 A review of the behavior of radioiodine in the subsurface at two DOE sites *Sci. Total Environ.* **691** 466–75
- Nevin K P and Lovley D R 2000 Potential for nonenzymatic reduction of Fe^{III} via electron shuttling in subsurface sediments *Environ. Sci. Technol.* **34** 2472–8
- Nevin K P and Lovley D R 2002 Mechanisms for accessing insoluble Fe^{III} oxide during dissimilatory Fe^{III} reduction by *Geothrix fermentans* *Appl. Environ. Microbiol.* **68** 2294–9
- Newcomer M *et al* 2021 Hysteresis patterns of watershed nitrogen retention and loss over the past 50 years in United States hydrological basins *Glob. Biogeochem. Cycles* **35** e2020GB006777
- Newsome L, Morris K, Shaw S, Trivedi D and Lloyd J R 2015 The stability of microbially reduced U^{IV}; impact of residual electron donor and sediment ageing *Chem. Geol.* **409** 125–35
- Nico P S, Anastasio C and Zasoski R J 2002 Rapid photo-oxidation of Mn(II) mediated by humic substances *Geochim. Cosmochim. Acta* **66** 4047–56
- Nico P S, Kumfer B M, Kennedy I M and Anastasio C 2009a Redox dynamics of mixed metal (Mn, Cr and Fe) ultrafine particles *Aerosol Sci. Technol.* **43** 60–70
- Nico P S, Stewart B D and Fendorf S 2009b Incorporation of oxidized uranium into Fe (hydr) oxides during Fe^{II} catalyzed remineralization *Environ. Sci. Technol.* **43** 7391–6
- Noël V *et al* 2017a Understanding controls on redox processes in floodplain sediments of the Upper Colorado River Basin *Sci. Total Environ.* **603** 663–75
- Noël V, Boye K, Kukkadapu R K, Li Q and Bargar J R 2019 Uranium storage mechanisms in wet-dry redox cycled sediments *Water Res.* **152** 251–63
- Noël V, Boye K, Lezama Pacheco J S, Bone S E, Janot N, Cardarelli E, Williams K H and Bargar J R 2017b Redox controls over the stability of U^{IV} in floodplains of the Upper Colorado River Basin *Environ. Sci. Technol.* **51** 10954–64
- O'Loughlin E J 2008 Effects of electron transfer mediators on the bioreduction of lepidocrocite (γ -FeOOH) by *Shewanella putrefaciens* CN32 *Environ. Sci. Technol.* **42** 6876–82
- O'Loughlin E J, Boyanov M I, Antonopoulos D A and Kemner K M 2011 Redox processes affecting the speciation of technetium, uranium, neptunium and plutonium in aquatic and terrestrial environments *Aquatic Redox Chemistry* (Washington, DC: ACS Publications) pp 477–517
- O'Loughlin E J, Boyanov M I, Flynn T M, Gorski C A, Hofmann S M, McCormick M L, Scherer M M and Kemner K M 2013 Effects of bound phosphate on the bioreduction of lepidocrocite (γ -FeOOH) and maghemite (γ -Fe₂O₃) and formation of secondary minerals *Environ. Sci. Technol.* **47** 9157–66
- O'Loughlin E J, Boyanov M I, Gorski C A, Scherer M M and Kemner K M 2021 Effects of Fe^{III} oxide mineralogy and phosphate on Fe^{II} secondary mineral formation during microbial iron reduction *Minerals* **11** 149
- O'Loughlin E J, Boyanov M I, Kemner K M and Thalhammer K O 2020 Reduction of Hg(II) by Fe^{II}-bearing smectite clay minerals *Minerals* **10** 1079
- O'Loughlin E J, Gorski C A, Flynn T M and Scherer M M 2019 Electron donor utilization and secondary mineral formation during the bioreduction of lepidocrocite by *Shewanella putrefaciens* CN32 *Minerals* **9** 434
- O'Loughlin E J, Kelly S D, Cook R E, Csencsits R and Kemner K M 2003a Reduction of uranium(VI) by mixed iron(II)/iron(III) hydroxide (green rust): formation of UO₂ nanoparticles *Environ. Sci. Technol.* **37** 721–7
- O'Loughlin E J, Kelly S D and Kemner K M 2010 XAFS investigation of the interactions of U^{VI} with secondary mineralization products from the bioreduction of Fe^{III} oxides *Environ. Sci. Technol.* **44** 1656–61
- O'Loughlin E J, Kemner K M and Burris D R 2003b Effects of Ag^I, Au^{III} and Cu^{II} on the reductive dechlorination of carbon tetrachloride by green rust *Environ. Sci. Technol.* **37** 2905–12
- O'Reilly A M, Chang N-B and Wanielist M P 2012 Cyclic biogeochemical processes and nitrogen fate beneath a

- subtropical stormwater infiltration basin *J. Contam. Hydrol.* **133** 53–75
- Olsen T A, Brandt C C and Brooks S C 2016 Periphyton biofilms influence net methylmercury production in an industrially contaminated system *Environ. Sci. Technol.* **50** 10843–50
- Olsen T A, Muller K A, Painter S L and Brooks S C 2018 Kinetics of methylmercury production revisited *Environ. Sci. Technol.* **52** 2063–70
- Özgen-Xian I, Kesserwani G, Caviedes-Voullième D, Molins S, Xu Z, Dwivedi D, Moulton J D and Steefel C I 2020 Wavelet-based local mesh refinement for rainfall-runoff simulations *J. Hydroinform.* **22** 1059–77
- Paddison L 2016 The world's threatened rivers - in pictures (available at: www.theguardian.com/sustainable-business/gallery/2016/sep/22/worlds-threatened-rivers-pollution-industry-agriculture-in-pictures) (Accessed 15 March 2021)
- Painter S L 2018 Multiscale framework for modeling multicomponent reactive transport in stream corridors *Water Resour. Res.* **54** 7216–30
- Painter S L 2021 On the representation of hyporheic exchange in models for reactive transport in stream and river corridors *Front. Water* **2** 69
- Painter S L, Coon E T, Atchley A L, Berndt M, Garimella R, Moulton J D, Svyatskiy D and Wilson C J 2016a Integrated surface/subsurface permafrost thermal hydrology: model formulation and proof-of-concept simulations *Water Resour. Res.* **52** 6062–77
- Painter T H et al 2016b The Airborne Snow Observatory: fusion of scanning lidar, imaging spectrometer and physically-based modeling for mapping snow water equivalent and snow albedo *Remote Sens. Environ.* **184** 139–52
- Pan C, Jiao Y, Kersting A B and Zavarin M 2021 Plutonium redox transformation in the presence of iron, organic matter and hydroxyl radicals: kinetics and mechanistic insights *Environ. Sci. Technol.* **55** 1800–10
- Paniconi C and Putti M 2015 Physically based modeling in catchment hydrology at 50: survey and outlook *Water Resour. Res.* **51** 7090–129
- Parks J M et al 2013 The genetic basis for bacterial mercury methylation *Science* **339** 1332–5
- Patton N R, Lohse K A, Godsey S E, Crosby B T and Seyfried M S 2018 Predicting soil thickness on soil mantled hillslopes *Nat. Commun.* **9** 1–10
- Pawloski G, Wurtz J, and Drellack S 2009 The underground test area project of the Nevada test site: building confidence in groundwater flow and transport models at Pahute Mesa through focused characterization studies *Technical Report Lawrence Livermore National Lab. (LLNL), Livermore, CA (United States)*
- Peña J, Bargar J R and Sposito G 2015 Copper sorption by the edge surfaces of synthetic birnessite nanoparticles *Chem. Geol.* **396** 196–207
- Peretyazhko T, Zachara J M, Heald S M, Jeon B-H, Kukkadapu R K, Liu C, Moore D and Resch C T 2008 Heterogeneous reduction of Tc^{VII} by Fe^{II} at the solid–water interface *Geochim. Cosmochim. Acta* **72** 1521–39
- Peterson M J et al 2018 Mercury remediation technology development for lower east fork poplar creek—FY 2017 Progress Report *Technical Report Oak Ridge National Lab. (ORNL), Oak Ridge, TN (United States)*
- Phelps T, Fliermans C, Garland T, Piffner S and White D 1989 Methods for recovery of deep terrestrial subsurface sediments for microbiological studies *J. Microbiol. Methods* **9** 267–79
- Plathe K L, Lee S-W, Tebo B M, Bargar J R and Bernier-Latmani R 2013 Impact of microbial Mn oxidation on the remobilization of bio-reduced U^{IV} *Environ. Sci. Technol.* **47** 3606–13
- Podar M, Gilmour C C, Brandt C C, Soren A, Brown S D, Crable B R, Palumbo A V, Somenahally A C and Elias D A 2015 Global prevalence and distribution of genes and microorganisms involved in mercury methylation *Sci. Adv.* **1** e1500675
- Powell B A, Dai Z, Zavarin M, Zhao P and Kersting A B 2011 Stabilization of plutonium nano-colloids by epitaxial distortion on mineral surfaces *Environ. Sci. Technol.* **45** 2698–703
- Powell B A, Fjeld R A, Kaplan D I, Coates J T and Serkiz S M 2005 Adsorption and reduction by synthetic hematite and goethite *Environ. Sci. Technol.* **39** 2107–14
- Powell R M, Puls R W, Hightower S K and Sabatini D A 1995 Coupled iron corrosion and chromate reduction: mechanisms for subsurface remediation *Environ. Sci. Technol.* **29** 1913–22
- Prancevic J P and Kirchner J W 2019 Topographic controls on the extension and retraction of flowing streams *Geophys. Res. Lett.* **46** 2084–92
- Prein A F, Holland G J, Rasmussen R M, Clark M P and Tye M R 2016 Running dry: the US Southwest's drift into a drier climate state *Geophys. Res. Lett.* **43** 1272–9
- Prikryl J D, Jain A, Turner D R and Pabalan R T 2001 Uranium(VI) sorption behavior on silicate mineral mixtures *J. Contam. Hydrol.* **47** 241–53
- Pruess K 1987 TOUGH user's guide
- Pruess K 2004 The TOUGH codes—A family of simulation tools for multiphase flow and transport processes in permeable media *Vadose Zone J.* **3** 738–46
- Qian Y, Yin X, Lin H, Rao B, Brooks S C, Liang L and Gu B 2014 Why dissolved organic matter enhances photodegradation of methylmercury *Environ. Sci. Technol. Lett.* **1** 426–31
- Ragnarsdottir K et al 2000 Uranium behaviour in natural environments *Environmental Mineralogy: Microbial Interactions, Anthropogenic Influences, Contaminated Land and Waste Management* pp 245–89
- Rathore S S, Jan A, Coon E T and Painter S L 2021 On the reliability of parameter inferences in a multiscale model for transport in stream corridors *Water Resour. Res.* **57** e2020WR028908
- Reguera G, McCarthy K D, Mehta T, Nicoll J S, Tuominen M T and Lovley D R 2005 Extracellular electron transfer via microbial nanowires *Nature* **435** 1098–101
- Reichstein M et al 2019 Deep learning and process understanding for data-driven Earth system science *Nature* **566** 195–204
- Ren H, Song X, Fang Y, Hou Z J and Scheibe T D 2021 Machine learning analysis of hydrologic exchange flows and transit time distributions in a large regulated river *Front. Artif. Intell.* **4** 39
- Riley W, Maggi F, Kleber M, Torn M, Tang J, Dwivedi D and Guerry N 2014 Long residence times of rapidly decomposable soil organic matter: application of a multi-phase, multi-component and vertically resolved model (BAMS1) to soil carbon dynamics *Geosci. Model Dev.* **7** 1335–55
- Riscassi A, Miller C and Brooks S 2016 Seasonal and flow-driven dynamics of particulate and dissolved mercury and methylmercury in a stream impacted by an industrial mercury source *Environ. Toxicol. Chem.* **35** 1386–400
- Rodell M, Velicogna I and Famiglietti J S 2009 Satellite-based estimates of groundwater depletion in India *Nature* **460** 999–1002
- Roden E E, Kappler A, Bauer I, Jiang J, Paul A, Stoesser R, Konishi H and Xu H 2010 Extracellular electron transfer through microbial reduction of solid-phase humic substances *Nat. Geosci.* **3** 417–21
- Roden E E and Wetzel R G 1996 Organic carbon oxidation and suppression of methane production by microbial Fe^{III} oxide reduction in vegetated and unvegetated freshwater wetland sediments *Limnol. Oceanogr.* **41** 1733–48
- Rogers D B et al 2021 Modeling the impact of riparian hollows on river corridor nitrogen exports *Front. Water* **3** 7
- Royer R A, Burgos W D, Fisher A S, Jeon B-H, Unz R F and Dempsey B A 2002 Enhancement of hematite bioreduction by natural organic matter *Environ. Sci. Technol.* **36** 2897–904

- Sani R K, Peyton B M, Dohnalkova A and Amonette J E 2005 Reoxidation of reduced uranium with iron(III)(hydr) oxides under sulfate-reducing conditions *Environ. Sci. Technol.* **39** 2059–66
- Santelli C M, Webb S M, Dohnalkova A C and Hansel C M 2011 Diversity of Mn oxides produced by Mn(II)-oxidizing fungi *Geochim. Cosmochim. Acta* **75** 2762–76
- Santschi P H, Roberts K A and Guo L 2002 Organic nature of colloidal actinides transported in surface water environments *Environ. Sci. Technol.* **36** 3711–19
- Sassen D S, Hubbard S S, Bea S A, Chen J, Spycher N and Denham M E 2012 Reactive facies: an approach for parameterizing field-scale reactive transport models using geophysical methods *Water Resour. Res.* **48** W10526
- Scheibe T D and Chien Y-J 2003 An evaluation of conditioning data for solute transport prediction *Groundwater* **41** 128–41
- Scheibe T D, Chien Y-J and Radtke J S 2001 Use of quantitative models to design microbial transport experiments in a sandy aquifer *Groundwater* **39** 210–22
- Scheibe T D, Fang Y, Murray C J, Roden E E, Chen J, Chien Y-J, Brooks S C and Hubbard S S 2006 Transport and biogeochemical reaction of metals in a physically and chemically heterogeneous aquifer *Geosphere* **2** 220–35
- Scheibe T D, Hubbard S S, Onstott T C and DeFlaun M F 2011 Lessons learned from bacterial transport experiments at the South Oyster Site *Ground Water* **49** 745–63
- Scheibe T D, Murphy E M, Chen X, Rice A K, Carroll K C, Palmer B J, Tartakovsky A M, Battiatto I and Wood B D 2015a An analysis platform for multiscale hydrogeologic modeling with emphasis on hybrid multiscale methods *Groundwater* **53** 38–56
- Scheibe T D, Schuchardt K, Agarwal K, Chase J, Yang X, Palmer B J, Tartakovsky A M, Elsethagen T and Redden G 2015b Hybrid multiscale simulation of a mixing-controlled reaction *Adv. Water Resour.* **83** 228–39
- Schmidt M W et al 2011 Persistence of soil organic matter as an ecosystem property *Nature* **478** 49–56
- Schwartz G E, Olsen T A, Muller K A and Brooks S C 2019 Ecosystem controls on methylmercury production by periphyton biofilms in a contaminated stream: implications for predictive modeling *Environ. Toxicol. Chem.* **38** 2426–35
- Senko J M, Istok J D, Suflita J M and Krumholz L R 2002 In-situ evidence for uranium immobilization and remobilization *Environ. Sci. Technol.* **36** 1491–6
- Senko J M, Kelly S D, Dohnalkova A C, McDonough J T, Kemner K M and Burgos W D 2007 The effect of U^{VI} bioreduction kinetics on subsequent reoxidation of biogenic U^{IV} *Geochim. Cosmochim. Acta* **71** 4644–54
- Senko J M, Mohamed Y, Dewers T A and Krumholz L R 2005a Role for (Fe^{III}) minerals in nitrate-dependent microbial U^{IV} oxidation *Environ. Sci. Technol.* **39** 2529–36
- Senko J M, Suflita J M and Krumholz L R 2005b Geochemical controls on microbial nitrate-dependent U^{IV} oxidation *Geomicrobiol. J.* **22** 371–8
- Sharon I et al 2015 Accurate, multi-kb reads resolve complex populations and detect rare microorganisms *Genome Res.* **25** 534–43
- Sharp J O et al 2011 Uranium speciation and stability after reductive immobilization in aquifer sediments *Geochim. Cosmochim. Acta* **75** 6497–6510
- Shi L, Richardson D J, Wang Z, Kerisit S N, Rosso K M, Zachara J M and Fredrickson J K 2009 The roles of outer membrane cytochromes of *shewanella* and *geobacter* in extracellular electron transfer *Environ. Microbiol. Rep.* **1** 220–7
- Shuai P et al 2019 Dam operations and subsurface hydrogeology control dynamics of hydrologic exchange flows in a regulated river reach *Water Resour. Res.* **55** 2593–2612
- Simanova A A, Kwon K D, Bone S E, Bargar J R, Refson K, Spósito G and Peña J 2015 Probing the sorption reactivity of the edge surfaces in birnessite nanoparticles using nickel(II) *Geochim. Cosmochim. Acta* **164** 191–204
- Šimůnek J and van Genuchten M T 2008 Modeling nonequilibrium flow and transport processes using HYDRUS *Vadose Zone J.* **7** 782–97
- Šimůnek J, Van Genuchten M T and Šejna M 2006 The HYDRUS software package for simulating two- and three-dimensional movement of water, heat and multiple solutes in variably-saturated media *Tech. Manual Version 1* 241
- Šimůnek J, van Genuchten M T and Šejna M 2008 Development and applications of the HYDRUS and STANMOD software packages and related codes *Vadose Zone J.* **7** 587–600
- Singer D M, Chatman S M, Ilton E S, Rosso K M, Banfield J F and Waychunas G A 2012 U(VI) sorption and reduction kinetics on the magnetite (111) surface *Environ. Sci. Technol.* **46** 3821–30
- Sivaswamy V et al 2011 Multiple mechanisms of uranium immobilization by *Cellulomonas* sp. strain ES6 *Biotechnol. Bioeng.* **108** 264–76
- Skylberg U, Bloom P R, Qian J, Lin C-M and Bleam W F 2006 Complexation of mercury(II) in soil organic matter: EXAFS evidence for linear two-coordination with reduced sulfur groups *Environ. Sci. Technol.* **40** 4174–80
- Smith D, Finnegan D and Bowen S 2003 An inventory of long-lived radionuclides residual from underground nuclear testing at the Nevada test site, 1951–1992 *J. Environ. Radioact.* **67** 35–51
- Song X, Chen X, Stegen J, Hammond G, Song H-S, Dai H, Graham E and Zachara J M 2018 Drought conditions maximize the impact of high-frequency flow variations on thermal regimes and biogeochemical function in the hyporheic zone *Water Resour. Res.* **54** 7361–82
- Sonnenthal E, Spycher N, Xu T and Zheng L 2021 TOUGHREACT V4. 12-OMP and TReactMech V1. 0 Geochemical and Reactive-Transport User Guide
- Spiro T G, Bargar J R, Spósito G and Tebo B M 2010 Bacteriogenic manganese oxides *Acc. Chem. Res.* **43** 2–9
- Spycher N F, Issarangkun M, Stewart B D, Şengör S S, Belding E, Ginn T R, Peyton B M and Sani R K 2011 Biogenic uraninite precipitation and its reoxidation by Fe^{III} (hydr) oxides: A reaction modeling approach *Geochim. Cosmochim. Acta* **75** 4426–40
- Steefel C I 2019 Reactive transport at the crossroads *Rev. Mineral. Geochem.* **85** 1–26
- Steefel C I, Carroll S, Zhao P and Roberts S 2003 Cesium migration in Hanford sediment: a multisite cation exchange model based on laboratory transport experiments *J. Contam. Hydrol.* **67** 219–46
- Steefel C I, DePaolo D J and Lichtner P C 2005 Reactive transport modeling: an essential tool and a new research approach for the Earth sciences *Earth Planet. Sci. Lett.* **240** 539–58
- Steefel C et al 2015 Reactive transport codes for subsurface environmental simulation *Computat. Geosci.* **19** 445–78
- Stegen J C and Goldman A E 2018 WHONDRS: a community resource for studying dynamic river corridors
- Stetten L et al 2018 Geochemical control on the reduction of U^{VI} to mononuclear U^{IV} species in lacustrine sediments *Geochim. Cosmochim. Acta* **222** 171–86
- Stewart B D, Girardot C, Spycher N, Sani R K and Peyton B M 2013 Influence of chelating agents on biogenic uraninite reoxidation by Fe^{III} (hydr) oxides *Environ. Sci. Technol.* **47** 364–71
- Stoliker D L et al 2013 Evaluating chemical extraction techniques for the determination of uranium oxidation state in reduced aquifer sediments *Environ. Sci. Technol.* **47** 9225–32
- Stucker V K, Silverman D R, Williams K H, Sharp J O and Ranville J F 2014 Thioarsenic species associated with increased arsenic release during biostimulated subsurface sulfate reduction *Environ. Sci. Technol.* **48** 13367–75
- Stucker V K, Williams K H, Robbins M J and Ranville J F 2013 Arsenic geochemistry in a biostimulated aquifer: an aqueous speciation study *Environ. Toxicol. Chem.* **32** 1216–23
- Stylo M, Alessi D S, Shao P P, Lezama-Pacheco J S, Bargar J R and Bernier-Latmani R 2013 Biogeochemical controls on the

- product of microbial U^{VI} reduction *Environ. Sci. Technol.* **47** 12351–8
- Suzuki Y, Kelly S D, Kemner K M and Banfield J F 2002 Nanometre-size products of uranium bioreduction *Nature* **419** 134–134
- Szecsody J E, Krupka K M, Williams M D, Cantrell K J, Resch C T and Fruchter J S 1998 Uranium mobility during *in situ* redox manipulation of the 100 Areas of the Hanford Site *Technical Report* Pacific Northwest National Lab. (PNNL), Richland, WA (United States)
- Taillefert M, Beckler J S, Carey E, Burns J L, Fennessey C M and DiChristina T J 2007 *Shewanella putrefaciens* produces an Fe^{III} -solubilizing organic ligand during anaerobic respiration on insoluble Fe^{III} oxides *J. Inorg. Biochem.* **101** 1760–7
- Tan H, Zhang G, Heaney P J, Webb S M and Burgos W D 2010 Characterization of manganese oxide precipitates from Appalachian coal mine drainage treatment systems *Appl. Geochem.* **25** 389–99
- Tang G, Watson D B, Wu W-M, Schadt C W, Parker J C and Brooks S C 2013a U^{VI} bioreduction with emulsified vegetable oil as the electron donor—model application to a field test *Environ. Sci. Technol.* **47** 3218–25
- Tang G, Wu W-M, Watson D B, Parker J C, Schadt C W, Shi X and Brooks S C 2013b U^{VI} bioreduction with emulsified vegetable oil as the electron donor—microcosm tests and model development *Environ. Sci. Technol.* **47** 3209–17
- Tang Y, Webb S M, Estes E R and Hansel C M 2014 Chromium(III) oxidation by biogenic manganese oxides with varying structural ripening *Environ. Sci.: Process. Impacts* **16** 2127–36
- Tokunaga T K, Wan J, Kim Y, Sutton S R, Newville M, Lanzirrotti A and Rao W 2008 Real-time x-ray absorption spectroscopy of uranium, iron and manganese in contaminated sediments during bioreduction *Environ. Sci. Technol.* **42** 2839–44
- Tolar B B, Boye K, Bobb C, Maher K, Bargar J R and Francis C A 2020 Stability of floodplain subsurface microbial communities through seasonal hydrological and geochemical cycles *Front. Earth Sci.* **8** 338
- Tripathi A, Tripathi D K, Chauhan D and Kumar N 2016 Chromium(VI)-induced phytotoxicity in river catchment agriculture: evidence from physiological, biochemical and anatomical alterations in *Cucumis sativus* (L.) used as model species *Chem. Ecol.* **32** 12–33
- Ulrich K-U, Ilton E S, Veeramani H, Sharp J O, Bernier-Latmani R, Schofield E J, Bargar J R and Giammar D E 2009 Comparative dissolution kinetics of biogenic and chemogenic uraninite under oxidizing conditions in the presence of carbonate *Geochim. Cosmochim. Acta* **73** 6065–83
- U.S. DOE 2019 *Open Watershed Science by Design: Leveraging Distributed Research Networks to Understand Watershed Systems Workshop Report* DOE/SC-0200, U.S. Department of Energy Office of Science (ess.science.energy.gov/open-watershed-workshop/)
- van Genuchten C M and Peña J 2016 Sorption selectivity of birnessite particle edges: a d-pdf analysis of Cd(II) and Pb(II) sorption by δ - MnO_2 and ferrihydrite *Environ. Sci.: Process. Impacts* **18** 1030–41
- Varadharajan C et al 2017 Reoxidation of chromium(III) products formed under different biogeochemical regimes *Environ. Sci. Technol.* **51** 4918–27
- Varol M and Şen B 2012 Assessment of nutrient and heavy metal contamination in surface water and sediments of the upper Tigris River, Turkey *Catena* **92** 1–10
- Vaughn L J, Conrad M E, Bill M and Torn M S 2016 Isotopic insights into methane production, oxidation and emissions in Arctic polygon tundra *Glob. Change Biol.* **22** 3487–502
- Veeramani H et al 2011 Products of abiotic U^{VI} reduction by biogenic magnetite and vivianite *Geochim. Cosmochim. Acta* **75** 2512–28
- Vörösmarty C J et al 2010 Global threats to human water security and river biodiversity *Nature* **467** 555–61
- Wainwright H M et al 2015 Identifying multiscale zonation and assessing the relative importance of polygon geomorphology on carbon fluxes in an Arctic tundra ecosystem *J. Geophys. Res.: Biogeosci.* **120** 788–808
- Wainwright H M et al 2020 Satellite-derived foreshore drought sensitivity of plant productivity in Rocky Mountain headwater catchments: spatial heterogeneity and geological-geomorphological control *Environ. Res. Lett.* **15** 084018
- Wainwright H M et al 2022 Watershed zonation through hillslope clustering for tractably quantifying above-and below-ground watershed heterogeneity and functions *Hydrol. Earth Syst. Sci.* **26** 429–44
- Wainwright H M, Orozco A F, Bucker M, Dafflon B, Chen J, Hubbard S S and Williams K H 2016b Hierarchical Bayesian method for mapping biogeochemical hot spots using induced polarization imaging *Water Resour. Res.* **52** 533–51
- Wainwright H et al 2016a Effective long-term monitoring strategies by integrating reactive transport models with *in situ* geochemical measurements 16212 WM2016 Conf pp 1–15
- Wainwright H, Sassen D, Chen J and Hubbard S 2014 Bayesian hierarchical approach for estimation of reactive facies over plume-scales using geophysical datasets *Water Resour. Res.* **50** 4564–84
- Wall J D and Krumholz L R 2006 Uranium reduction *Annu. Rev. Microbiol.* **60** 149–66
- Wan J, Tokunaga T K, Brodie E, Wang Z, Zheng Z, Herman D, Hazen T C, Firestone M K and Sutton S R 2005 Reoxidation of bioreduced uranium under reducing conditions *Environ. Sci. Technol.* **39** 6162–9
- Wang S, Jaffe P R, Li G, Wang S and Rabitz H A 2003 Simulating bioremediation of uranium-contaminated aquifers; uncertainty assessment of model parameters *J. Contam. Hydrol.* **64** 283–307
- Wang Z, Lee S-W, Catalano J G, Lezama-Pacheco J S, Bargar J R, Tebo B M and Giammar D E 2013a Adsorption of uranium(VI) to manganese oxides: x-ray absorption spectroscopy and surface complexation modeling *Environ. Sci. Technol.* **47** 850–8
- Wang Z, Lee S-W, Kapoor P, Tebo B M and Giammar D E 2013b Uraninite oxidation and dissolution induced by manganese oxide: a redox reaction between two insoluble minerals *Geochim. Cosmochim. Acta* **100** 24–40
- Wang Z, Tebo B M and Giammar D E 2014a Effects of Mn^{II} on UO_2 dissolution under anoxic and oxic conditions *Environ. Sci. Technol.* **48** 5546–54
- Wang Z, Ulrich K-U, Pan C and Giammar D E 2015 Measurement and modeling of U^{IV} adsorption to metal oxide minerals *Environ. Sci. Technol. Lett.* **2** 227–32
- Wang Z, Xiong W, Tebo B M and Giammar D E 2014b Oxidative UO_2 dissolution induced by soluble Mn(III) *Environ. Sci. Technol.* **48** 289–98
- Watson D B et al 2013 *In situ* bioremediation of uranium with emulsified vegetable oil as the electron donor *Environ. Sci. Technol.* **47** 6440–8
- Weinan E, Bjorn E and Zhongyi H 2003 Heterogeneous multiscale method: a general methodology for multiscale modeling *Phys. Rev. B* **67** 092101
- Wentz D A, Brigham M E, Chasar L C, Lutz M A and Krabbenhoft D P 2014 Mercury in the nation's streams—levels, trends, and implications *Technical Report* US Geological Survey
- White M D, Bacon D H, McGrail B P, Watson D J, White S K and Zhang Z 2012 STOMP subsurface transport over multiple phases: STOMP-CO₂ and STOMP-CO₂e guide: version 1.0 *Technical Report* Pacific Northwest National Lab. (PNNL), Richland, WA (United States)
- White M D and Oostrom M 2000 STOMP subsurface transport over multiple phases version 2.0 theory guide *Technical Report* Pacific Northwest National Lab. (PNNL), Richland, WA (United States)

- White M D and Oostrom M 2003 STOMP subsurface transport over multiple phases version 3.0 User's guide *Technical Report* Pacific Northwest National Lab., Richland, WA (US)
- Wiatrowski H A, Das S, Kukkadapu R, Ilton E S, Barkay T and Yee N 2009 Reduction of Hg^{I} to Hg^0 by magnetite *Environ. Sci. Technol.* **43** 5307–13
- Wilkins M J et al 2009 Proteogenomic monitoring of *Geobacter* physiology during stimulated uranium bioremediation *Appl. Environ. Microbiol.* **75** 6591–9
- Williams K H et al 2011 Acetate availability and its influence on sustainable bioremediation of uranium-contaminated groundwater *Geomicrobiol. J.* **28** 519–39
- Williams K H, Bargar J R, Lloyd J R and Lovley D R 2013 Bioremediation of uranium-contaminated groundwater: a systems approach to subsurface biogeochemistry *Curr. Opin. Biotechnol.* **24** 489–97
- Wrighton K C et al 2012 Fermentation, hydrogen and sulfur metabolism in multiple uncultivated bacterial phyla *Science* **337** 1661–5
- Wu R, Chen X, Hammond G, Bisht G, Song X, Huang M, Niu G-Y and Ferre T 2021 Coupling surface flow with high-performance subsurface reactive flow and transport code PFLOTTRAN *Environ. Modell. Softw.* **137** 104959
- Wu W-M et al 2006a Pilot-scale *in situ* bioremediation of uranium in a highly contaminated aquifer. 1. Conditioning of a treatment zone *Environ. Sci. Technol.* **40** 3978–85
- Wu W-M et al 2006b Pilot-scale *in situ* bioremediation of uranium in a highly contaminated aquifer. 2. Reduction of U^{VI} and geochemical control of U^{VI} bioavailability *Environ. Sci. Technol.* **40** 3986–95
- Wu W-M et al 2010 Effects of nitrate on the stability of uranium in a bioreduced region of the subsurface *Environ. Sci. Technol.* **44** 5104–11
- Xu C, Santschi P, Roberts K, Zhong J, Hatcher P, Hung C, Francis A, Dodge C and Honeyman B 2008 Colloidal cutin-like siderophoric molecules mobilize plutonium from contaminated soils of the Rocky Flats Environmental Technology Site (RFETS), USA *Environ. Sci. Technol.* **42** 8211–17
- Xu S and Jaffé P R 2006 Effects of plants on the removal of hexavalent chromium in wetland sediments *J. Environ. Qual.* **35** 334–41
- Yabusaki S B et al 2007 Uranium removal from groundwater via *in situ* biostimulation: field-scale modeling of transport and biological processes *J. Contam. Hydrol.* **93** 216–35
- Yabusaki S B et al 2011 Variably saturated flow and multicomponent biogeochemical reactive transport modeling of a uranium bioremediation field experiment *J. Contam. Hydrol.* **126** 271–90
- Yabusaki S B et al 2017 Water table dynamics and biogeochemical cycling in a shallow, variably-saturated floodplain *Environ. Sci. Technol.* **51** 3307–17
- Yabusaki S B, Steefel C I and Wood B 1998 Multidimensional, multicomponent, subsurface reactive transport in nonuniform velocity fields: code verification using an advective reactive streamtube approach *J. Contam. Hydrol.* **30** 299–331
- Yan J, Wang L, Hu Y, Tsang Y F, Zhang Y, Wu J, Fu X and Sun Y 2018 Plant litter composition selects different soil microbial structures and in turn drives different litter decomposition pattern and soil carbon sequestration capability *Geoderma* **319** 194–203
- Zachara J M et al 2013 Persistence of uranium groundwater plumes: contrasting mechanisms at two DOE sites in the groundwater–river interaction zone *J. Contam. Hydrol.* **147** 45–72
- Zachara J M, Chen X, Song X, Shuai P, Murray C and Resch C T 2020 Kilometer-scale hydrologic exchange flows in a gravel bed river corridor and their implications to solute migration *Water Resour. Res.* **56** e2019WR025258
- Zachara J M, Fredrickson J K, Li S-M, Kennedy D W, Smith S C and Gassman P L 1998 Bacterial reduction of crystalline Fe^{+3} oxides in single phase suspensions and subsurface materials *Am. Mineral.* **83** 1426–43
- Zachara J M, Girvin D C, Schmidt R L and Resch C T 1987 Chromate adsorption on amorphous iron oxyhydroxide in the presence of major groundwater ions *Environ. Sci. Technol.* **21** 589–94
- Zachara J M, Smith S C, Liu C, McKinley J P, Serne R J and Gassman P L 2002 Sorption of Cs^+ to micaceous subsurface sediments from the Hanford site, USA *Geochim. Cosmochim. Acta* **66** 193–211
- Zachara J, Cowan C, Schmidt R and Ainsworth C 1988 Chromate adsorption by kaolinite *Clays Clay Miner.* **36** 317–26
- Zachara J, Fredrickson J, Li S, Kennedy D, Smith S and Gassman P 1995 Bacterial reduction of crystalline Fe^{3+} oxides in single phase suspensions and subsurface materials *Am. Mineral.* **83** 11–12:1426–1443
- Zaunbrecher L K, Cygan R T and Elliott W C 2015 Molecular models of cesium and rubidium adsorption on weathered micaceous minerals *J. Phys. Chem. A* **119** 5691–700
- Zhang J, Ma T, Yan Y, Xie X, Abass O K, Liu C, Zhao Z and Wang Z 2018 Effects of Fe-S-As coupled redox processes on arsenic mobilization in shallow aquifers of Datong Basin, northern China *Environ. Pollut.* **237** 28–38
- Zhao P, Zavarin M, Dai Z and Kersting A B 2020 Stability of plutonium oxide nanoparticles in the presence of montmorillonite and implications for colloid facilitated transport *Appl. Geochem.* **122** 104725
- Zhao P, Zavarin M, Leif R N, Powell B A, Singleton M J, Lindvall R E and Kersting A B 2011 Mobilization of actinides by dissolved organic compounds at the Nevada Test Site *Appl. Geochem.* **26** 308–18
- Zheng Z, Tokunaga T K and Wan J 2003 Influence of calcium carbonate on U(VI) sorption to soils *Environ. Sci. Technol.* **37** 5603–8
- Zhuang K, Izallalen M, Mouser P, Richter H, Risso C, Mahadevan R and Lovley D R 2011 Genome-scale dynamic modeling of the competition between *Rhodospirillum rubrum* and *Geobacter* in anoxic subsurface environments *ISME J.* **5** 305–16
- Zyvoloski G A, Robinson B A, Dash Z V and Trease L L 1997a Summary of the models and methods for the FEHM application—a finite-element heat-and mass-transfer code *Technical Report* Los Alamos National Lab., NM (US)
- Zyvoloski G A, Robinson B A, Dash Z V and Trease L L 1997b User's manual for the FEHM application—A finite-element heat-and mass-transfer code *Technical Report* Los Alamos National Lab